1480 e.7

Tractatus, (cui Titulus, *Proceſſus Integri in Morbis ferè Omnibus Curandis*) à Clariſſimo D. D. *Thomâ Sydenham* Conſcriptus, Imprimatur.

*Thomas Burwell*, Præſes.

*Humphrey Brook*,
*Joſ. Clerk*,
*Guilelmus Briggs*,
*Fredr. Slare*,
} Cenſores.

Datum 18° Die *Auguſti* A. D. 1693. in Comitiis Cenſoriis ex Ædibus Collegii noſtri.

# PROCESSUS INTEGRI
## IN
# MORBIS
Ferè Omnibus
# CURANDIS,
## A
Dº THOMA SYDENHAM, M. D.
Conſcripti.

Quibus acceſſit graphica Symptomatum delineatio, unà cum quamplurimis Obſervatu dignis.

NECNON

*De* PHTHISI *Tractatulo.*

EDITIO QUARTA.

*LONDINI:*
Impenſis *J. Knapton* ad Inſigne Coronæ, & *Guil. Innys* ad Inſignia Principis, in Cœmeterio D. Pauli. MDCCXII.

22
SEP
58

# AMICE LECTOR,

SPECIMEN habes Praxeos eximiæ, in dilectissimi filii M. D. usum, cautè & diligenter excogitatum, suâ ipsius manu conscriptum, mihi autem pro familiaritate istâ quâ omnes excepit concessum à *Cl. Sydenham*, Viro sanè, cui vita superior parem dederat, vel postera daturæ est, non modò in arte suâ mirâ sagacitate, & ingenii acumine singulari, conspicuo: sed & probis moribus, & in humanum genus, studio & amore insignibus adeò, ut apud omnes summum sui reliquerit desiderium: Cujus præcepta licèt animo meo & imo pectore infixa sint, nè tamen fortè fortunâ in mei ipsius & ægrorum detrimentum excidant memoriâ, earùm fidâ, typis haud ita pridem mandabam, exemplaribus circiter viginti impressis, ut amicis quoque inservirem. Nescio autem quo pacto in lucem irrepsit publicam inter *Miscellanea Curiosa* superioris anni Norimbergæ editum;

tum; ſed ineptâ mole inutile. Prodiit tandem eo modo veſtitum ut tibi ſit in quovis loco paratum; figmentis cerebri vanis & futilibus ſciolorum dogmatis nequaquam refertum, nec hypotheſi placidâ ſolidis ut videtur rationum momentis undequaque conſtipatâ ſuperbiens; humile autem & modeſtum accuratè morbos depingit, & accuratiùs adhuc medetur; in id ſolum ſollicitum, ut innoteſcat quod natura faciat aut ferat. Et ſi verum loqui liceat, modò partium ſitus noveris, pateant tibi per Symptomata ægritudines, (quod aſſiduâ ediſcas obſervatione) ſcias deindè veram & genuinam medendi methodum per diætam & Pharmaciam adhibere, fæliciſſimus eris practicus, & tibi & infirmis optimè conſules. Parùm interim tuâ refert utrum Alcali peccet an Acidum, num in Spiritibus Animalibus, an in ipſo ſanguine, aut in recondito aliquo Viſcere deliteſcat morbi minera; unde de Periodico Intermittentium reditu doctè ſatis & prolixè poſſis diſſerere; dum interea ipſa Febris ſine ullis ambagibus, inquietudine, ſiti, calore, pulſu celeri, nauſeâ, cæteriſque Symptomatis, etiam aſtantibus palam eſt, & apertè dignoſcitur: Hinc ſæpius mirari ſubit, quamobrem anxiè ita & ſcrupulosè immediatas & proximas morborum cauſas, dum Medicinam pertractandam ſuſcipiunt, indagare ſatagunt, & è tenebris iſtis, quibus natura obduxit, irrito licèt conatu, eruere, gravi & ſubacto judicio

viri

iri & in praxi verſatiſſimi; ſatis interim gna- ſe latere vel quævis obvia. Undè enim di- nanat graminis viridis, aut albus nivis color? Quid eſt quòd humana anima ratiocinari ne- queat dum adhuc Infantes aut Pueri ſumus? Undè forma tua Specifica? Satius eſſet Medi- cum quàm Philoſophum agere (quis enim un- quam Carteſium ſibi Medicum voluerit?) mi- nutiſſima Phœnomena planè & ritè enume- rare, & remedia quibus unuſquiſque morbus faciliùs cedat, candido & ſincero animo pro- ponere; ita tandem limites ſuos prætergreſſa, quibus anguſtis nimis jam includitur ars no- biliſſima, omnibus, quaquaverſum patet orbis, fæliciter ſatis & proſperè ſalutem, filiis autem ſuis honores ſummos conciliabit.

Meditabatur paulò ante obitum, tractatum de Phthiſi peritiſſimus medicus; in perquiren- do naturæ methodum, quâ morbos amoliri ſtudet, ut adjutrices porrigeret manus, & mi- ſeriis hominum ſubveniret, diu noctuque pen- ſitabundo, & anxio animo, exercitatiſſimus; hinc fidiſſimarum obſervationum ſerie & irre- fragabili edoctus experientiâ, medicinæ praxin ornavit, & ad ſaniores reduxit regulas; ſed væ tabidis, dum hujuſmodi nunquam intermiſſis ſtudiis detenti & defatigati ſpiritus, ſolitis non vacabant excubiis, truculentiſſimus hoſtis, *Podagra*, quæ per multos annos obſederat, & de- bilitaverat artus, impetu facto in interiora vi- ſcera irruit, per vomitum & ſeceſſum enor-

miores

miores intentans mortem: Unico autem morbo neſcit cedere Medicorum Princeps, ſed cum malorum cumulo acceſſit & *Rerum Calculus*, qui corraſis vaſorum oſculis, ſanguinem liberaliùs, quàm ſenectus tulit, urinæ permixtum effudit, ſe jam non ampliùs opponens, nec vitæ, nec mortis cupidus, placidam, & pacatam, expiravit animam. Nè tamen auxilio prorſùs diſtituantur tabidi fragmenta quædam, quæ autorem ubivis ſapiunt, in primo ſaltem morbi ſtadio plerumque proficua, (neque enim fas eſt, quod ægrotanti proximo opituletur, celare,) in publicum prodeunt; ut exindè pateat quàm accurata, & omnibus numeris abſoluta, ſi modo Deus Opt. Maximus conceſſiſſet vitam, expectari potuiſſet Phthiſiologia.

Quid in Febribus, Variolis, Morbillis, acutis cæteris, ut & Chronicis ſanandis valuit optimè meritus autor noſter, ſequentes indicant paginæ. Regimen & Diæta inſtituuntur aptiſſima; præſcribuntur remedia pauca, ſed neceſſaria; nec in Pharmacopolarum lucrum excogitata, nec inani oſtentationi compoſita: ſitibundis ægris (eodem modo quo & ipſe ſitim ſuam reſtinguere ſolitus erat) cereviſiam tenuem affatim & pro libitu hauriendam, undè mirè reficiuntur & refrigerantur, concedit; nec immanitate plùs quàm barbarâ, precibus eorum ſurdus, Apozemata & Julapia præ ingrato ſapore faſtidita ſubſtituit; maximam adhibet

libet cautelam ne exæſtuans nimis, vel igne culentiori, vel ſtragulorum onere, plùs adhuc calefceret, vel à propinatis forſan remediis um in finem, ut per poros eliminetur mate-ia morbifica cruda adhuc & inconcocta, quæ amen hâc datâ ansâ concitato motu cere-brum petit, undè Phreneſis aut Coma, aut ex-travaſato ſanguine deturpant cutim maculæ purpureæ, aut eruptiones, quas vocant milia-res, collum & pectus ubivis obſident. Nun-quid ad amuſſim magis depingi poterat quàm Variolarum hiſtoria? eruptionis diem in utra-vis ſpecie deſignat, puſtularum ingenium ad unguem deſcribit, quo tempore incipiat, & quamdiu continuetur Salivatio, quo iterum fa-ciei manuumque tumor ſe prodat; quid tan-dem quolibet die expectandum, diſertè explic-cat. Paregoricorum uſum, quo emolumento judicent Practici, primus perſuaſit, malum, ingerendi Cardiaca ante eruptionem, morem, undè ſæpè accerſitum fuerit confluentium ge-nus, meritò primus improbavit. Meliùs au-tem hæc ex ipſo fonte. Valeas itaque amice Lector, & ſicubi ſcateant Mendæ ignoſcas.

*S. M.*

Amico

## Amico ſuo Integerrimo M. D

### Expertiſſimo Domino *S. M.*

*Solum ſequaces pratulimus diu*
*Coum Medentes; Compar habebitur*
*Qui clariorem te, minori*
*Reddidit hoc, Medicina! Libro.*

*Non hic tumenti Mole ſuperbiens*
*Blattis futurus Præda opulentior:*
*Senſu ſed arcta Ditiori*
*Conciliant ſibi Verba Pondus.*

*Fruſtrà moleſto vaſtu Volumine*
*Luſtres Galeni ſcripta; Minutula*
*Acumen huic majus Tabellæ;*
*Se ſtupet hic Medicina totum.*

*Hic non remoto ſordidus Angulo*
*Senſim fatiſcet; Sedula ſtrenuè*
*Manus quotannis, Spe perenni*
*Deteret, nitidior ut reſurgat.*

*Applaude Votis Turba Medentium:*
*Non Febris illum, Morbus atrocior*
*Haud afflet ullus; Commodis qui*
*Immoritur ſatagendo Veſtris.*

Sic vovet,

*Chriſtophorus Crelle*, M. D.

# INDEX.


A

DE Abortu præcavendo Pag. 63
Ambustis 85
Anginâ 30
Apoplexiâ 45
Arthmate 89

C

Canis rabidi morsu 88
Chlorosi 82
Cholerâ Morbo 55
Chorea sancti Viti 43
Colicâ biliosâ 58
Colicâ Hystericâ 60
Colicâ Pictonum, 56
Contusionibus 87

D

Diabete 76
Diarrhœâ 50
Dysenteriâ ibid.

E

Epilepsiâ Puerorum 77

F

Febre Stationariâ, 19
Febre Constitutionum variarum 15
Febribus à dentitione ortis 80
Febre depuratoriâ 12
Febre Erysipelatosâ 28
Febre Hecticâ Infantum 80
Febre Intermitt. 16
Febre Pestilentiali 14
Febre Scarlatinâ 22
Fluore albo 74
Furfure sive Porrigine capitis 87

G

Gonorrhœâ virulentâ 69

H

Hæmoptoë 83
Hæmorrhagiâ narium 81
Hæmorrhoidum dolore 76
Hæmorrhoidum fluxu immodico 77


Hy-

Hydrope p. 65
Hypochondriacâ affectione 5
Hyſtericâ Paſſione ib.

I

Ictero 62
Iliacâ Paſſione 57

L

Lochiorum fluxu immodico 63
Lochiorum ſuppreſſione 64
Lue Venereâ 72

M

Maniâ communi 85
Menſium fluxu immodico 58
Menſium ſuppreſſione 83
Mictu Sanguineo à Renum calculo 49
Morbillis 31

N

Nephritico Paroxyſmo 48

O

Ophthalmiâ 47

P

Paralyſi 90
Peripneumoniâ nothâ 24
Phthiſi 91, 94
Pleuritide 23
Pruritu ſerino 29
Punctûrâ tendinis 85
Podagrâ 100

R

Rachitide 79
Rheumatiſmo 25
Remediorum formulæ 1

S

Scorbuto 93

T

Tuſſi 91
Teneſmo 50
Tuſſi Puerorum convulſivâ 81

V

Variolis 34
Ulcere Veſicæ 88
Vomitu ſanguinis 83
Uteri procidentiâ 48

*Medi-*

# *Medicamentorum quædam formulæ in praxi magis familiares.*

## Potio Communis Purgativa.

℞ *Tamarind.* ℥ſs. *fol. Sennæ* ℥ij. *Rhei* ℥iſs. *coq. ſ. q. aq. ad* ℥iij. *in Colatura diſſolve Mannæ & Syrup. Roſar. ſolut. an.* ℥j, *m. f. Potio.*

## Emeticum Commune.

℞ *Aq. Cardui Benedict.* ℥ij. *Infuſ. Croc. Metal.* ℥j. *Syr. Caryophyll.* ℥ſs. *m. f. vomitorium.* Sumat horâ 4ª Pomeridianâ, ſuperbibendo poſt ſingulas vomitiones hauſtum largum liquoris poſſetici.

## Julapium Perlatum.

℞ *Aq. Ceraſor. nigr. lact. Alexeter. an.* ℥iij. *aq. Cinnamom. hordeat.* ℥j. *Margarit. præparat.* ℥iſs, *ſacchar. Cryſtall. q. ſ. adde aq. Roſar. Damaſcen.* ℥ſs. *f. Julap.* de quo cap. cochlear. *iv*, vel *v*, in Languoribus.

## Julapium Cardiacum.

℞ *Aq. Lact. Alexeter. Ceraſor. nigr. an.* ℥iij. *aq. Epidem. Syr. Caryophyll. & de ſucco Citri, an.* ℥ſs, *m. f. Julap. de quo capiat ſapiùs.*

## Decoctum Diæteticum.

℞ *Rad. Sarſaparill.* ℥vj. *lign. Saſſafr. & Chinæ ana* ℥ij. *Glycyrrhiz.* ℥j. *coq. in aq. font. Congiis ij, per horæ dimidium; deindè ſtent clausè ſuper cineres calidos per horas xij, poſteà ebulliant ad tertiæ partis conſumptionem; Cum primùm ab igne amoveatur, infunde ſem. Aniſi.* ℥ſs. *poſt duas horas coletur, & poſtea depuretur per reſidentiam, & liquor reponatur lagenis vitreis probè clauſis ad uſum.* Sit pro potu ordinario per *xxx* dies.

## Apozema Aperiens & Antiſcorbuticum.

℞ *Rad. gramin. Cichor. Fœnicul. & Aſparag. ana* ℥j. *uvarum Corinthiac. & paſſul. enucleat. an.* ℥ij. *fol. hepat. ling. cerv. & cap. Ven. an. M. j. fol. Beccabung. ſub finem addit. M. ij; coq. ſ. q. aq. fontan. ad* ℔ij. *ſub finem adde vini Rhenani* ℔ſs. *Colaturâ adhuc calente, infunde ad duas horas, vaſe clauſo, Cochl. hortenſ. M. j. colatur. adde ſyr. è quinque radic. & è ſucco Aurant. ana* ℥ij. *aq. Cinnamom. hordeat.* ℥j. *M. f. Apozem.* Sumat ℔ſs. manè & 5ª pomeridianâ, per *xiv* dies.

Eclegma

## Eclegma Incraſſans in Tuſſi.

℞ *Ol. Amygd. dulc.* ℥j. *Syr. Papav. Rhœad. Portulac. Jujub. & Lohoch. ſani, ana* ℥ſs. *Sacchar. cryſtal. q. ſ. M. in mortario marmoreo ad integræ horæ ſpatium, & f. Linctus perfectè miſtus; recondatur in ollâ figulinâ, ad uſum.* Cap. ſæpiùs lambend. è baculo Liquiritiæ.

## Eclegma fortiùs incraſſans.

℞ *Conſerv. Roſ. rub. Syr. violar. & de Mecon. an.* ℥j. *ſem. papav. alb.* ℥iij. *contundantur ſimul & per ſetaceum trajiciantur, deindè adde ol. nuc. moſchat. expreſſ. gr. vj. f. Eclegma.*

## In Defluxione tenui.

℞ *Conſerv. Roſ. rub.* ℥ij. *Syr. de Mecon. & jujub. an.* ℥j. *Oliban. Maſtich. & Succin. an.* ℥ſs. *ol. nuc. Moſchat. per expreſſionem gr. vj. M. f. Lohoch,* de quo lambat ſæpè; bis in die cap. eclegm. deſcript. cochl. j, inſtillando Bilſam. Sulphur. Aniſat. à gut. *viij,* ad gut. *xij.*

## Cereviſia Purgativa.

℞ *Polypod. Querc.* ℔ſs. *Rad. Rhabarb. Monach. fol. ſenna & paſſul. enucleat. ana* ℔ſs. *Rhabarb. inciſ. & rad. Raphan. Ruſtican. an.* ℥iij. *fol. Cochl. hortenſ. ſalv. an. M. iv, Aurantior. inciſ.* N° iv, *infunde in Congiis v, vel vj, Cereviſia non lupulata, fermentationis tempore,*

*& quando ad maturitatem pervenerit,* ſit pro potu ordinario per dies *xiv,* vel *xxj,* præcipuè verò ſumat hauſtum ſingulis Auroris.

## Emplaſtrum Hyſtericum.

℞ *Galbani in tinctura Caſtor. ſolut. & colat.* ℥iij. *Tacamahac.* ℥ij. *f. Emplaſt. pro umbilico.*

## Catharticum pro Infantibus tenellis.

℞ *Syrup. Cichor. cum Rhabarb. Cochl. j. parvulum.*

## Decoctum Amarum Purgativum.

℞ *Decoct. amar. duplicatâ ſenâ* ℥iv. *Syrup. de Spinâ Cervinâ* ℥j. *Elect. de Succ. Roſar.* ℥ij. *M. f. Potio.*

## Laudanum Liquidum Authoris.

℞ *Vini Hiſpanici* ℔j. *Opii* ℥ij. *Croci* ℥j. *pulv. Cinnamomi, & Garyophillorum, an.* ℥j. *infundantur ſimul in B. M. per duos vel tres dies, donec liquor debitam conſiſtentiam acquirat; Colatura ſervetur pro uſu.*

DE

# DE AFFECTIONE,

## IN

## *Fœminis, hyſtericâ; maribus, hypochondriacâ dictâ.*

CUM à caſu aliquo graviori moleſtum patitur humana anima, ataxiam concipiunt animales ſpiritus; urina ſubindè limpidiſſima copiosè excernitur; ſanitatis recuperandæ ſpes omnis protinùs abjicitur, & triſtiſſima quævis ſibi ominantur ægrotæ; quamcunque corporis partem morbus inſederit (plurimas autem infidet) ſymptomata ei parti competentia confeſtim urgent; in capite apoplexia, hemiplegiâ ſoluta, ſtatim à partu adoritur; ſpaſmi epilepſiæ perſimiles (vulgò uteri ſtrangulatus, in quo venter & præcordia guttur versùs intumeſcunt)

infeſtant; clavus hyſtericus, in quo dolor in capite atrociſſimus, quem tranſverſo pollice poſſis contegere, ægrâ interim materiam viridem bili porraceæ parem evomente, miſerè excruciat; cordis palpitationem, tuſſim, paſſiones iliacam & colicam, nephritin, urinæ ſuppreſſionem æmulatur paroxyſmus; vomitiones enormes, diarrhœam ſubindè excitat; externè in carne muſculosâ nunc dolorem, nunc tumorem inducit in tibiis hydropis inſtar; nec (quod mireris) intactos dimittit dentes; dolor dorſi creberrimè ſævit; & ſæpè ſæpiùs adeò refrigerantur externæ partes, ut cadaver planè referant; in riſum & lachrymas effuſiſſimas ſinè ullâ παρφάσει ridiculè erumpunt; ptyaliſmus nonnunquam eo gradu laborantes afficit, ut Mercurio delibutas facilè crederes. Dolores hyſterici, quamcunque partem obſident, ubi diſceſſerint, teneritudinem, quæ tangi retuſat, relinquunt, ac ſi verberatæ fuiſſent carnes.

Mittatur Sanguis ad ℥viij.

*Emplaſt. de Galbano* umbilico admoveatur. Sequenti manè aggrediatur uſum ſequentium pilularum;

℞ *Pil. Coch. maj.* ℥ij. *Caſtor. pulv. gr. ij. Balſam. Peruv. gutt. iij. f. Pil. xij.* Cap. *iv.* ſingulis vel alternis matutinis, pro ratione virium, horâ 4ª vel 5ª matutinâ, ſuperdormiendo.

℞ *Aq.*

℞ *Aq. Ruta* ℥iv. *Bryon. comp.* ℥ij. *Caſtor. nodulo ligat. & in vitro ſuſpenſ.* ℥ſs. *Sacchar. Cryſtal. q. ſ. f. julap.* de quo capiat cochl. *iv*, vel *v*, in omni languore. Poſt has pilulas ſumptas utatur ſequentibus;

℞ *Limatur. Chalyb. gr. viij. cum ſ. q. Extract. Abſinth. f. Pil.* N° *iij.* Sumat ſummo mane, & repetantur horâ 5ª pomeridianâ per *xxx* dies, ſuperbibendo hauſtum Vini Abſinthiris.

Sin forma boli magis arrideat,

℞ *Conſerv. Abſinth. Roman. & Conſerv. fluved. Aurant. an.* ℥j. *Angelic. condit. Nuc. Moſchat. condit. & Theriach. Andromach. an.* ℥ſs. *pulv. Ari comp.* ℥iij. *Zinzib. condit.* ℥ij. *cum ſ. q. Syrup. de ſucc. citr. vel, ejus defectu, Syr. Aurantior. f. Electuarium.*

℞ *Elect. præſcript.* ℥ij. *limatur. Chalyb. gr. viij. cum ſ. q. Syrup. Aurantior. f. Bolus* ſumendus mane & ſerò, ſuperbibendo hauſtum vini abſinthitis, vel Cochl. *vj*, ſeq. Infuſionis.

℞ *Rad. Angelic. Enul. campan. & Imperator. an.* ℥j. *fol. Abſinth. vulg. Centaur. min. Marrub. alb. & Chamædr. an M. j. Cort. Aurant. duor. inciſ. affunde vini Hiſpanic. q. s. ad ſupereminentiam duorum digitorum.* Coletur tempore uſus. Vel,

In Delicatulis *Chalybs* exhiberi poterit in formâ pulv. ſequentis;

℞ *Limatur. Chalyb. probè trit.* ℥j, *pulv. Ari comp.* ℥vj. *ſem. Coriand. præparat. Aniſi, Fœnicul. dulc. an.* ℥ſs. *Cinnamom. acutiſſ. Corall. rubr. præparat. an.* ℥iij. *Nuc. Moſchat.* ℥ij. *f. omnium pulvis ſubtiliſſimus, cui adde ſacchar. albiſſ. ad pondus omnium,* hujus cap. è Cochleari ℥ſs. bis in die per 4 dies, tum ℥j. bis in die per 40 dies, ſuperbibend. Cochl. *vi,* ſeq. Julap. vel vini Abſinthitis.

℞ *Aq. Lact. Alexeter.* ℥xij. *Gentian. comp.* ℥iv. *Abſinth. magis comp.* ℥ij. *ſacchar. albiſſ. q. ſ. f. Julap.* Vel;

℞ *Vini Rhenan. Abſinth.* ℏſs. *aq. Gentian. comp.* ℥ij. *Syr. Caryophyll.* ℥j. *f. Julap.*

℞ *Myrrh. elect. Galban. & Aſæ fœtidæ, an.* ℥j. *Caſtor.* ℥ſs. *cum ſ. q. Balſam. Peruv. f. Pil. è ſingulis drachmis xij.* Sumat *iij,* ſingulis noctibus, ſuperbibend. aq. Bryon. compoſ. Cochl. *iij,* vel *iv,* per totum decurſum hujus proceſſûs.

Si Pilulæ Præſcriptæ ventrem moverint, utatur ſequentibus;

℞ *Caſtor.* ℥j. *ſal. ſuccin. volatil.* ℥ſs. *cum ſ. q. extract. Ruta, f. pilula* N° *xxiv.* Cap. *iij,* ſingulis

ſingulis noctibus, ſuperbibendo Julapii hyſteric. Cochl. *iij*, vel *iv*.

*Sp. C. C.* ad gutt. *xvj*, vel *xviij*, in aquâ aliquâ appropriatâ ſæpiùs exhibitus valde confert.

Si verò his non cedat remediis, tum capiat ſequentes pilulas;

℞ *Trochiſc. de Myrrh. pulv.* ℈*j. Balſam. ſulphur. Terebinth. gutt. iv. cum ſ. q. gumm. Ammoniac. ſolut. f. Pil.* N° *iv.* Cap. manè & ſerò ſuperbib. Julap. hyſteric. Cochl. *iv*, vel *v*, inſtillando Sp. C. C. gutt. *xij*.

*Electuarium Antiſcorbutic.* cum aquâ ibidem deſcriptâ prodeſt in his affectionibus, ut & *Electuar. corroborans cum additione Conſerv. Cochl. hortenſ.* ℥*j. pulv. Ari Compoſ.* ℥*vj*. ſuperbibendo præſcriptam aquam.

At ſi affectus his non cedat remediis, eundum eſt ad *aquas ferreas*; & ſi neque his, tum ad *Sulphureas*, quales ſunt *Bathonienſes*.

Inter bibendum aquas ferreas hæc obſervanda; ſi aliquod Symptoma gravius acciderit, quod potationi aquarum referri poteſt, tunc ab iis æger abſtineat, donec Symptoma iſtud penitùs evanuerit. Æger in earum uſu per ſex ſeptimanas ad minimum, vel potiùs, ad duos menſes perſeveret. *Zinziber conditum*, vel *Semina Carui Saccharata*, ad calefaciendum ventriculum ſubindè ſumat. *Pilulæ* etiam *hyſtericæ iij*. devorari poſſunt per primas *x* noctes, ſuperbibendo *Julap. hyſteric.* cochl. *iv*, vel *v*.

Quoàd uſum *aquarum Bathonienſium*, duobus diebus bibat eas, ac tertio die per modum Balnei eas ingrediatur, atque ita alternatim per ſex ſeptimanas vel duos menſes.

Si *Chalybs* juſto plùs corpus calefaciat, tum per ejus uſum, quartâ quâque Aurorâ ſumat *aq. mineralium purgantium* ℔iv. Quæ tametſi alvum moveant, haud tamen tumultum cient, ut ſolent Cathartica ex officinis deprompta.

Si *Chalybs* magnas perturbationes inducat, Laudanum aq. aliquâ hyſtericâ miſtum, ſingulis noctibus ad tempus aliquod propinandum.

Cùm à morbi diuturnitate fractæ & attritæ fuerint vires, venæſectio, & cathartica non ſunt præmittenda, ſed à Chalybe ſtatim incipiendum.

Ubi Symptomata ſunt mitiora, ſatis eſt ſanguinem mittere, & alvum ſolvere ad dies *iij*, vel *iv*, dein pilulas hyſtericas ad dies *x*, manè & ſerò exhibere.

In dolore intolerabili, vomitione, & *Diarrhœâ* enormioribus, exhibendum Laudanum, & poſtea corroborandi ſpiritus, at modò per vires liceat, antè uſum Laudani, educatur è venis ſanguis, & Catharticum propinetur; in viraginibus præſertim, & ſanguineo temperamento præditis, in debilioribus autem, & haud ita pridem paroxyſmum paſſis, ſufficit ventriculum Congio uno lactis cereviſiati hauſto, & mox per vomitum rejecto detergere, dein largam doſin *Theriac. Androm.* vel *Elect. Orvietan.*

*tam.* imperare, & Cochl. aliquot, liquoris alicujus Spirituosi, cum pauculis Laudani liquid. guttulis, superbibenda, dare. Quòd si ægra jampridem vomuerit, & ulteriùs emeticis sollicitare periculo non vacabit, Laudanum sinè morâ propinandum est, eâ dosi, quæ Symptoma vincat, & post singulas vomitiones toties quoties repetendum, in formâ potissimùm solidâ, vel in liquidâ, in parco vehiculo, viz. gutt. aliquot in cochl. *j.* aq. Cinamom. fortior. ægrâ interim conquiescente, & caput immotum tenente.

Perdomito Symptomate, manè & serò, ad dies pauculos continuetur usus Laudani.

Hæc duo diligentur notanda; quòd cum semel post evacuationes Laudano uti coeperimus, eâ dosi, atque illâ toties repetitâ, sumendum est, donec Symptoma omninò fuerit debellatum, interjecto tantùm spatiolo inter singulas doses, ut quid prior præstiterit ante noscamus quàm alteram ingeramus, deinceps cum Laudano morbum aggredimur, nihil movendum est, nihil evacuandum vel per mitissimum enema.

*Theriaca Andromachi* crebrò, diuque usurpata, magnum est in hoc, & aliis morbis quàm plurimis à coloris, & concoctionis defectu ortis, remedium.

Vina *Hispanica* gentianâ, angelicâ, absinth. centaurio, corticum aurantior. flavedine, aliisque corroborantibus infusis medicata, si ad aliquot

quot Cochlearia ter in die potentur, admodum conferunt, modo ægra tenui non fuerit biliosoque corporis habitu.

*Cortex. Peruv.* ad ℈j. manè & serò, per septimanas aliquot deglutitus, præsertim in spasmis hystericis miras habet vires.

In tenui biliosoque corporis habitu possit usurpari Diæta lactea, præcipuè in Colicâ Hystericâ, si modò ea incommoda possint superari, quæ primis ferè diebus apportat, quod nempè tum in ventriculo coaguletur, tum etiam viribus corporis, in statu conservandis, sustendandisque minùs sufficiat.

Nihil tamen æquè sanguinem spirituſque fovet, & confirmat, ac diu multumque singulis ferè diebus equo vehi; nec itinera in curru facta suo carent successu.

*De Febre Depuratoriâ annorum 1661, 62, 63, 64.*

Si æger juvenis sit, extrahatur sanguis è brachio, & eodem die post aliquot horas, vel sequenti die duabus horis post leve prandium, *Emeticum ex infus. Croc. Metall. paratis Zythogal.* ℔ *vj. vel viij.* ut quoties vomuerit, aut alvum exoneraverit, è vestigio sumatur haustus, exhibeatur; & post finitam operationem Emetici, hoc vel simile paregoricum;

℞ *Aq.*

℞ *Aq. Ceraſor. nigr.* ℥*iſs. aq. Epidem.* ℥*ſs. Laudan. liquid. gutt. xvi. Syr. Caryophyll.* ℥*ij. M. F. Hauſtus.*

Vomitoria ex infuſ. *Croc. Metall.* parata, puerulis, ulliſve infra adoleſcentiam conſtitutis, vel minimâ quantitate, non eſt tutum exhibere.

Poſt hunc diem uſque ad undecimum vel duodecimum ſingulis Auroris injiciatur enema ſequens;

℞ *Decoct. commun. pro Clyſtere* ℔*j. (vel lactis vaccini eandem quantitatem) Sacchar. Culinar. & Syrup. violac. an.* ℥*ij. f. Enema.*

Poſt dictum diem alvus conſtipari permittatur, quo materia febrilis citiùs concoquatur, ad quem etiam finem Cardiaca moderata his ultimis diebus exhibita apprimè conducunt; ut,

℞ *Pulv. è chel. Cancror. comp. gr. xiv. Electi. de Ovo* ℈*ſs. cum ſ. q. Syr. Caryophyll. f. Bolus* ſumendus octavâ quâque horâ, ſuperbibendo ſequentis Julapii Cochl. *v*, vel *vj*.

℞ *Aq. Lact. Alexeter. Ceraſor. nigr. an.* ℥*iiij. Aq. Epidem. & Syr. Caryophyll. an.* ℥*j. M. F. Julap.* vel alia his ſimilia.

Hac methodo ſollicitè obſervatâ, uſitatò circa *xv* diem (tum ex laudabili in urina ſeparatione, tum ex manifeſta Symptomatum om-

omnium remiſſione) tempeſtivum eſſe Potionem Purgantem communem exhibere percipies.

Accidit interdum, maximè in Senibus, curatâ Febre, & purgato corpore, ægrum nihilominus valdè debilem eſſe, & quandoque tuſſi, interdum etiam ſcreatu, magnam glutinoſi viſcoſique phlegmatis copiam expectorare. Hoc in caſu vinum Malaganum annoſum, cum pane toſto ei immiſſo, bibat æger.

Si ſuperveniat Iliaca paſſio, præſcribatur Sal. Abſinth. ℈j. ex Cochl. j. Suc. Limonum, manè & ſerò, intermediis temporibus cap. Aq. menth. ſtillatit. ſine Saccharo, Cochl. aliquot, bis quâque horâ. Dum hæc fiunt, catulus vivens nudo ventri indeſinenter accumbat. Poſtquam dolor cum vomitu, per biduum triduumve ceſſaverit, exhibeatur *Pil. Coch. maj.* in *Aq. Menth.* ſoluta, nec removendus catulus ante pilularum uſum.

Ad recidivam præcavendam, perſiſtat in uſu prædictæ aquæ per longum tempus, & duplicatis pannis laneis ventrem à frigore defendat.

## *De Febre Peſtilentiali 1665, 66.*

Poſt venæſectionem in lecto celebrandam, æger ſtragulis obruatur, & Lacinia lanea ſinciputi alligetur; tum, ſi vomitus non adſit, hoc vel ſimile Hydroticum exhibe.

℞ *The-*

℞ *Theriac. Andromach.* ℥ſs. *Elect. de ov.* ℈j. *Pulv. è chel. cancr. comp. gr. xij, Coccinel gr. viij. Croc. gr. iv. cum ſ. q. ſucc. kermes, f. Bolus*, quem ſumat ſextâ quâque horâ, ſuperbibendo Cochl. *vj*, ſeq. Julapii.

℞ *Aq. Card. Benedict. & Scord. compoſ. an.* ℥iv. *Aq. Theriacal. ſtillatit.* ℥ij. *Syr. Caryophyl.* ℥j. *M. F. Julapium.*

Si adſit vomitio, hydroticum differatur, donec ſolo tegumentorum pondere ſudor promanare incipiat, linteaminis parte vultui ſuperinductâ.

Incepta Diaphoreſis hauſtulis Zythogalæ ſalviâ alteratæ, vel Cereviſiæ cui macis aliquantulum incoxeris ſubinde repetitis ad naturalis diei ſpatium (i. e. horas 24.) eſt continuanda; in ſudore licet juſculis confortativis ægrum reficere.

Ubi tumor enatus fuerit, venam aperire non ſuſtinui; per horas adhuc à finito ſudore *xxiv*, ægro ſe in lecto continente frigus cautè devitetur. Induſium ſponte ſuâ in corpore areſcere permittatur, potulenta omnia calidiuſculè hauriantur, & Zythogalæ ſalviâ alteratæ uſus etiamnum continuetur. In ſeq. luce potio purgat. communis exhibeatur.

Anni 1667, 1668. Febrim Varioloſam & Variolas produxerunt.

Anni 1669, 70, 71, 72. Febrim Dyſentericam.

Anni

Anni 1673, 74, 75. Febrim Comatoſam. Curationem febrium horum annorum non attingo; quoniam methodo in febre anni 1685, &c. deſcriptâ curari eas potuiſſe autumo.

*De Febribus Intermittentibus.*

Cum rigore & horrore adoritur paroxyſmus, quibus mox ſuccedit calor, & huic tandem ſudor, & ſubſequens ἀπυρεξία: licèt in primis morbi diebus, Autumno maximè, aliquoties remiſſio potius quàm intermiſſio perſentiſcitur. In calore æque ac in rigore vomiturit utplurimum ac vehementer, ægrotat miſer cum ſiti & lingua arida. In tenera ætate abdominis inflatio, in adultis crurum tumor febris ſolutionem; tonſillarum dolor, raucedo, oculi cavi, facies Hippocratica, mortem denunciant.

℞ *Cort. Peruv. ſubtiliſſ. pulv.* ℥j. *cum ſ. q. Syr. Caryophyll. vel de Roſis ſiccis, f. Electuarium dividendum in xij doſes*; quarum unam capiat quartâ quâque horâ, ſuperbibendo Vini cujuſlibet hauſtum, incipiend. immediate poſt paroxyſmum.

Si autem præſcripti boli ventrem moverint, in hauſtu Vini dentur *gutt. x Laudan. liquid.* ſingulis vel alternis vicibus poſt aſſumptionem boli, ut opus fuerit.

Ad

Ad recidivam præcavendam (præcipuè in quartana) repetatur proceſſus prædictus ſingulis ſeptimanis per tres alias vices.

Sin forma Pilularum magis arrideat,

℞ *Cort. Peruv. pulv.* ℥*j. cum ſ. q. Syrup. caryophyll. f. Pilula mediocris magnitudinis*, quarum capiat *vj*, quartâ quâque horâ.

℞ *Cort. Peruv. pulv.* ℥*ij. Vin. Rhenan.* ℔*ij. ſtent ſimul in Infuſione frigida, & coletur per manicam* Hippocratis. Uſûs tempore cap. ℥*iij.* tertiâ vel quartâ quâque horâ.

Sin ob vomituritionem, ferè continuam, Corticem deglutire nequeat æger, tum ſexies, vel octies, Spatio duarum horarum, *Succi Limonum* recens expreſſi Cochl. *j*, cum *Salis Abſinth.* ℈*j.* aſſumat, dein *Laudan. Liquid. gutt. xvj*, cum *Aq. cinnam.* fortior. Cochl. *j*, ac brevi poſt, ceſſante vomitu, Corticis uſum aggrediatur.

In intermittentibus vernalibus, nonnunquam emeticum, tempeſtivè propinatum, ut nempè ante paroxyſmum negotio ſuo defungi poſſit, fœliciſſime ſucceſſit. Nonnunquam Enema in diebus intermittentibus, per triduum, vel quatriduum injectum, Curationem abſolvit. Quæ ſequenti quoque modo tentari poteſt.

℞ *Serpentar. Virgin. ſubtiliſſ. pulveriſat. gr. xv. Vin. alb.* ℥*iij.* Sumat duabus horis ante

Paroxyſ-

Paroyſmum, & ſtragulis coopertus æger ſudet per *iij*, vel *iv* horas, idem faciendo per duas alias vices ante Paroxyſmum.

Si ab ingeminatis paroxyſmis ſumme debilis fuerit æger;

℞ *Conſerv. Flor. Borag. Bugloſſ. an.* ℥*j. conſerv. Anthos.* ℥*ſs. cort. citr. condit. Nuc. Moſchat. condit. Theriac. Andromach. an.* ℥*iij. confection. Alchermes* ℥*ij. M. F. Opiata*; cap. ad quantitatem avellanæ, mane & veſperi, ſuperbibendo Cochlearia paucula, alicujus *Aq. Epidemicæ* ſimplicioris, *Saccharo* edulcoratæ, interdicto interim enematum uſu.

Si ſub morbi exitum occurrat hydrops, nondum tamen depulsâ Febre, curatio non eſt inſtituenda remediis purgantibus, ſed infuſionibus rad. *Raphan. Ruſtican. Summitat. Abſinth. centaur. Min. Baccar. Juniper. cinerum geniſt.* &c. in vino factis, exulante autem Febre, Cathartics, & aperientibus.

Pro Infantibus Febre intermittente laborantibus:

℞ *Aq. ceraſor. nigr. & Vini Rhenan. an.* ℥*ij. cort. Peruv. ſubtiliſſ. pulv.* ℥*ij. Syr. caryophyll.* ℥*j. M. F. Julap.* Cap. Cochl. *j.* alterumve, pro ratione ætatis quartâ quâque horâ, donec Paroxyſmi non amplius diſcrucient, inſtillando alternis vicibus, ſi Diarrhœa adfuerit, Laud. liquid. gutt. *j*, vel *ij*.

De

*De Febre Stationaria, ab an.* 1685, *ad* 90.

Calor & frigus ſe invicem excipiunt, artus pariter & Caput dolent, pulſus interim fere ſanus, nonnunquam tuſſis, dolor circa collum & fauces, exacerbatio Febris prope noctem, inquietudo, ſitis, lingua vel humida & ex omni parte pelliculâ albâ & ſcabrâ veſtita, vel ſicca, fuſci coloris in medio, quam partem undique cingit fimbria inalbeſcens. A jugi in lecto decubitu coma & phreneſis, à calido regimine petechiæ, maculæ purpureæ, eruptiones (quas vocant) miliares morbillis magis rubentes, pulſus inordinatus, tendinum ſubſultus, atque ita tandem mors. Sudores in initio ſymptomatici divexant, qui ſi ab arte provocentur, viſcoſi à Capite erumpunt, & materiam morbificam in Caput aut in artus tranſferunt.

Mittat Sanguis ad ℥x è brachio. Reiteretur Venæſectio, ſi modò reſpirationis difficultas, dolor Capitis inter tuſſiendum lancinans, & alia ſigna Peripneumoniæ nothæ adſint. In hoc caſu tam venæſectio quàm Catharſis repetenda, donec æger convaluerit.

Veſperi applicetur Epiſpaſticum, & ſequenti Aurorâ Potio Lenitiva præſcribatur, quæ ad tertiam vicem alternis diebus repetatur, & horâ ſomni poſt Catharſin hauſtus pacificus propinetur.

℞ *Aq.*

℞ *Aq. Paralyſ.* ℥iij. *Syr. de Mecon.* ℥ſ. *Succ. Limon. recent. extract. cochl. ij. M. F. Hauſtus.*

Aphthæ & ſingultus ſponte ſuâ poſt curationem febris ſupervenientes ſponte evaneſcunt; Si in longum tempus decurrant, facilè fugantur *Cortic. Peruv.* ℥ſ. *cum Syr. Papav. Rhœad. q. ſ. in formam Electuar. vel pilular.* redactâ, ſuperbibend. hauſtum lactis deflorati. Remedium hoc certiſſimè ſcopum attinget, ſi non fruſtrentur ejus vires ex jugi ægri in lecto decubitu.

Diebus à purgatione liberis ſequentia præſcribantur.

℞ *Conſerv. Luſul. & cynosbat. an.* ℥ſ. *conſerv. Berber.* ℥ſſ. *crem. Tart.* ℥ſ. *cum ſ. q. Syr. de Limonib. f. Electuarium*, de quo cap. ad magnitud. Nuc. Moſchat. ter in die, ſuperbibendo ſequent. Julap. Cochl. *vj.*

℞ *Aq. Portulac. Lactuc. Paralyſ. an.* ℥iij. *Syr. de Limonib.* ℥iſſ. *Syr. Violar.* ℥ſ. *M. f. Julapium.* Vel,

℞ *Aq. font.* ℔ſ. *aq. Roſar. Succ. Limonum, Sacchar. albiſſ. an.* ℥iv. *coq. lento igne dum deſpumaverint, cap.* ℥iij. *ad libitum.* præſcribatur quoque ſeq. gargariſma.

℞ *Succ. Pomor. agreſt.* ℔ſſ. *Syr. de rubo Idæo* ℥ſ. *m. f. gargariſma.*

Si

Si febris ita ventriculum infeſtet, ut æger potionem retinere nequeat, *Pil. Coch. maj. ad* ℈*ij.* exhibeantur, tum etiam hypnoticum eſt concedendum, *v. gr. Laudan. Londin. gr. iſs. cum ſimili quantitate Maſtich.* vel *Laudan. Liquid. gut. xviij, in aq. Cinnamom. hordeat.* ℥*j.*

*Decoctum album paratum ex* ℥*j.* C. C. *uſti, in aq. communis* ℔*iij. cocti, & poſtea Saccharo albiſſimo edulcoratum* ſit pro potu, ut & Cereviſia tenuis pro libitu.

Poſt Catharſin bis celebratam, concedatur ægro, Pullorum carnibus veſci, & poſt ultimam Purgationem, modo febris aliquantulum deflagraverit, vini Canarini cochl. *iij*, vel *iv*, manè, & ſero, ut & poſt Prandium, per dies aliquot propinentur.

In Phrenitide, ut & Comate, nihil perinde confert, atque ægri caput radi, nullo admoto Emplaſtro, ſed Pileolo tantum.

Si (quod fœminis hyſtericis quandoque accidit) poſt venæſectionem, & purgationes, huiuſmodi iteratas, etiamnum duret, procurratque febris, modo ſigna Peripneumoniæ, & inflammationis circa partes vitales abſint, ſinguis noctibus Paregoricum, & medicamenta hyſterica, bis vel ter in die, injungantur.

## *De Infantibus Febre Stationariâ laborantibus.*

*Hirudines duæ* utrinque ponè aures applicentur, dein *Emplaſtr. Epiſpaſticum* cervici admoveatur.

Ce-

*Cerevisiâ, quâ Rhabarbarum fuerit infusum, purgetur.*

Si post Catharsin febris intermittere videatur, tum *Julapium cum Cortice Peruviano* (in Capite de febre intermittenti pro infantib. descriptum) exhibeatur.

*De Febre Scarlatina.*

Æstate exeunte, infantes præcipuè infestat. Rigent & horrent sub initium; nec tamen admodùm ægrotant; cutis universa maculis parvis, rubris, crebrioribus, latioribus & magis rubentibus quàm in morbillis, interstinguitur. Ad duos treve dies persistunt, quo quidem tempore evanescunt, & decedente subjectâ cuticulâ squammulis furfuraceis, ad instar farinæ, cutis conspergitur.

℞ C. C. *usti & pulv. è Chel. cancr. comp. an.* ℥ss. *Coccinel. gr. ij, Sacchar. Crystal.* ℥j. *M. f. pulvis subtilissᵗ. in xij Chartulas dividend.* quarum unam cap. 6. quâque horâ, superbibendo cochl. *ij*, vel *iij*, sequentis Julapii.

℞ *Aq. Ceraſor. nigr. aq. Lactis Alexeter. an.* ℥iij. *Syr. è succ. citri* ℥j. *M. f. Julap.*

Applicetur etiam Epispasticum cervici, & qualibet nocte *paregoricum ex Syr. de Mecon.* propinetur, & postquam symptomata cessaverint, purgetur æger.

Di

## *De Pleuritide.*

Inter Ver & Æſtatem dominatur. A rigore & horrore orditur, quos ſtatim comitantur calor, ſitis, inquietudo & cætera febrium ſymptomata; elapſis pauculis horis, æger vehementi dolore punctorio in laterum alterutro circa coſtas corripitur, qui nunc verſùs omoplatas, nunc ſpinam, nunc anteriora pectoris ſeſe propagat, tuſſi crebrâ excruciatur. Materia ſputo rejecta in initio morbi pauca eſt & tenuis, & ſæpe ſanguinis particulis intermiſta, progreſſu verò copioſior & magis concocta, ſanguine etiam permixto. Febris pari paſſu incedit cum tuſſi, ſputo ſanguinis, dolore, & pro ratione liberioris expectorationis ſenſim imminuitur; nunc alvus eſt aſtricta, nunc ſoluta nimis. Eductus è vena ſanguis ſevi liquati ſpeciem præ ſe fert, ubi refrixerit.

Mittatur ſanguis è brachio lateris affecti ad ℥x.

℞ *Aq. Papav. Rhœad.* ℥iv. *Sal. Prunel.* ℥j. *Syr. Violar.* ℥j. *M. f. Hauſtus*, ſumendus ſtatim à prima venæſectione.

℞ *Amygd. dulc. excorticat.* N° v, *ſem. Melon. Pepon. an.* ℥ſs. *ſem. papav. alb.* ℥iij. *aq. hord.* ℔iſs. *aq. Roſar.* ℥ij. *cum ſ. q. ſacchar. cand. f. ſ. a. Emulſio. Cap.* ℥iv. 4ª quâque horâ.

℞ *De-*

℞ *Decoct. Pectoral.* ℔ij. *Syr. Violar. & Cap. Ven. an.* ℥iſs. *M. f. Apozema. Cap.* ℔ſs. *ter in die.*

℞ *Ol. Amygd. dulc.* ℥ij. *Syr. Violar. & Cap. Ven. an.* ℥j. *Sacchar. Cand. q. ſ. M. f. Eclegma,* de quo ſæpe lambat.

*Oleum Amygd. dulc. per ſe,* vel *Oleum ſem. Lini recenter extract.* exhiberi poteſt.

℞ *Ol. Amygd. dulc. Lilior. & unguent. Dialthæa, an.* ℥j. *M. f. Linimentum,* quo inungatur latus affectum manè & ſerò, ſuperimponendo fol. Braſſicæ.

Repetatur Venæſectio ad tres alias vices, ad eandem quantitatem, quatuor diebus continuis, ubi dolor & reſpirandi difficultas ſuadent.

## *De Peripneumonia notha.*

Ingruente hyeme & ſæpe ſub ejus exitum emergit, inſultu primo nunc incaleſcit, nunc friget æger, ex motu quolibet vertiginoſus eſt, rubent & inflammantur genæ oculique, tuſſit, & inter tuſſiendum dolor capitis lancinans premit, potulenta vomitu rejicit, urina turbida eſt & intensè rubet, ſanguis pleuriticorum æmulus, anhelus crebrò & celeriter ſpiritum ducit, dolente thorace. Ab aſthmate ſicco planè diſtinguitur, quòd in iſto nuſquam comparent febris indicia, in hac manifeſtò ſe produnt,

dunt, licèt minora longè & magis obſcura quàm in peripneumonia vera.

Mittatur ſanguis è brachio dextro ad ℥x. Die ſequenti æger hanc potionem ſumat.

℞ *Caſſiæ extract.* ℥j. *Glycyr.* ℥ij. *ficus pingues* N° *iv. fol. ſenn.* ℥iiſs. *Agaric. trochiſcat.* ℥j. *coq. ſ. q. aquæ ad* ℥iv. *in Colatur. diſſolv. Mannæ* ℥j. *Syr. Roſar. ſolut.* ℥ſs.

Si verò potionem aſſumere nequeat, Cap. *Pil. coch. maj.* ℈ij. horâ quartâ matutinâ.

Sequenti die celebretur iterum venæſectio, ad eandem quantitatem; ac poſtero die repetatur Catharſis, quæ porrò repetenda eſt alternis diebus, vel poſt majora intervalla, pro ratione virium; Atque inſuper ſi Symptomata non ceſſaverint, extrahatur iterum ſanguis ſemel adhuc, bis, vel ſæpius, diebus aliquot interjectis, prout res poſtulaverit; ſed ut plurimum venæſectio bis celebrata ſufficiet. Interim, dum hæc fiunt, præſertim in diebus à purgatione vacuis, utatur *decocto Pectorali, Eclegmate* & *oleo Amygd.* dulcium, in Pleuritide deſcriptis.

## *De Rheumatiſmo.*

Rigorem & horrorem & cætera febrium ſymptomata patiuntur, elapſo die uno alterove (eſt & ubi citius) dolor adeſt atrox, nunc in hoc nunc in illo artu, in carpio, humeris &

genubus præſertim, viciſſim hos relinquens & illos occupans, rubore, & tumore in parte, quam poſtremùm affecit, adhuc reſiduis. Febris ſenſim evaneſcit, manente dolore, quin & nonnunquam immaniùs ſæviente. In lumbagine rheumatica immanis dolor adeſt, circa lumborum regionem fixus, nephritin mentiens, niſi quòd æger non vomituriat, qui procumbere nequit, ſed vel è lecto exilit, vel ſuper eodem erectus ſedet, corpore perpetuâ agitatione nunc versùs anteriora, nunc poſteriora inclinato. Sanguis pleuriticorum ſimilis.

Mittatur Sanguis è brachio lateris affecti, ad ℥x.

℞ *Aq. Nymph. Portulac. Lactuc. an.* ℥iv. *Syr. de Limonib.* ℥iſs. *Syr. Violac.* ℥j. *M. f. Julap.* de quo bibat ad libitum.

Præſcribend. *Emulſio 4. Sem. frigid. maj. Cataplaſma è mica panis albiſſ. & lacte Croco tincto* applicetur parti dolenti.

Die ſequenti eadem ſanguinis quantitas detrahatur; atque iterum, die uno vel altero interjecto, reiteretur, atque porrò quater vel pluries, ſi opus fuerit, hoc obſervando, ut poſt ſecundam Venæſectionem majora ſint intervalla inter ſingulas ſanguinis miſſiones.

Diebus à Phlebotomia vacuis Enema è lacte ſaccharato ſubinde injiciatur, vel ſequens;

℞ *De-*

℞ *Decoct. commun. pro Clyſtere* ℔j. *Syr. Violar. & Sacchar. culinar. an.* ℥ij. *M. f. Enema.*

Si Imbecillitas ægri repetitionem Phlebotomiæ non ferat, tum poſt ſecundam vel tertiam venæſectionem Curatio hac methodo tentanda.

Capiat alternis diebus *potionem purgativam communem*, donec convaluerit, & iiſdem noctibus Paregoric. *ex Syr. de Mecon.*

Sin his remediis non cedat, & æger ob imbecillitatem evacuationes qualeſcunque ferre nequeat, aggrediatur uſum *Elect. & aq. Antiſcorbutic.* in ſcorbuto deſcript. quæ etiam remedia locum habent in Rheumatiſmo Scorbutico.

In Juvenibus, atque iis qui temperatè vixerunt, vino non plus æquo dediti, Rheumatiſmus Diætâ ſimplici admodùm refrigerante, & mediocriter nutriente, æque fœliciter, ac repetitis Venæſectionibus (quas minus ferre queant) profligari poteſt. Ex. grat.

Solo Lactis ſero ad dies quatuor veſcatur æger, poſtea præter ſerum pane ſimilagineo ſemel in die loco prandii, donec penitus convaleſcat; niſi quòd poſtremis diebus iterum panem in cœnam indulgere liceat. Symptomatis ceſſantibus, carnes pullorum elixorum, cæteraque eupepta concedantur, tertio autem quoque die ſolo lactis ſero victitet, donec tandem pancraticè valeat.

*De Febre Eryſipelatoſa.*

Nulla non corporis pars, præſertim facies, tumet, dolet & ſummè rubet, puſtulæ denſæ, parvæ, in veſiculas ſubinde faceſſunt, quæ per frontem caputque ſerpunt, oculi tumore obruti, rigor & horror atque reliqua febris indicia adſunt. In alia hujus morbi ſpecie, liquorum attenuantium potationi ſuperveniente, febricula adeſt, puſtulæ urticarum puncturas referentes, nonnunquam in veſiculas ſublatæ, mox recedentes, tuberculorum more ſub cute ſe condentes, maximè pruriunt, & ſcalpendo rurſum apparent. Eſt & alia eruptionis ſpecies, pectus ut plurimum obſidens, macula nimirum lata, cuti vix ſupereminens, porriginoſa, cum ſquammulis flavis, qua vigente, æger bene ſe habet, eâdem evaneſcente, leviter ægrotat, urina turbida & flava, iiſdem remediis, quibus pruritus ferinus, cedit, vino utatur æger & carnibus eupeptis.

Mittatur ſanguis è brachio ad ℥*ix.* vel *x.* Die ſequenti ſumat *potionem communem purgativam.*

℞ *Rad. Althæa & Lilior. an.* ℥*j. fol. Malv. Sambuc. Verbaſc. flor. Chamæmel. Melilot. ſummitat. Hyperic. & centaur. minor. an. M. j. ſem. Lini, fœnugræc. an.* ℥*ſs. coq. ſ. q. aq. ad* ℔*iij. Coletur liquor, & tempore uſûs adde ſingulis*

*gulis libris decocti Sp. Vini* ℥*ij.* Stuphæ è panno laneo huic decocto immersæ, ac expressæ, calidè admoveantur bis in die, & pars affecta post fotum sequenti mixturâ illinatur.

℞ *Sp. Vini* ℔ß*s. Theriac. Androm.* ℥*ij. pulv. piperis longi, & Caryophyll. an.* ℥*ij. f. mixtura.* quâ Charta Emporetica madefacta parti affectæ circumvolvatur.

Si primæ venæsectioni non cedat morbus, tum repetatur; & nondum cessante, ad duas alias vices celebretur, interposito semper die uno.

Diebus à Venæsectione liberis *Enema è lacte cum Syr. Violarum, Emulsio refrigerans & Julap. refrigerans,* sunt præscribenda.

*De pruritu ferino, & inveteratis cutis eruptionibus, quæ Venæsectioni & Catharsi non cedunt.*

℞ *Theriac. Andromach.* ℥*ß. Elect. de ovo* ℥*j. rad. Serpentar. Virgin. subtiliss. pulver. gr. xv. Lapid. Bezoar. oriental. gr. v. cum s. q.* Syr. è *conditur. citri f. Bolus,* horâ somni sumendus per dies *xxj.* superbibendo cochl. *vj.* sequent. Julapii.

℞ *Aq. Card. Benedict.* ℥*vj. aq. Epidem. & Theriacal. stillat. an.* ℥*ij. Syrup. Caryophyll.* ℥*j. M. f. Julap.*

Singulis matutinis cap. liquoris poſſetici ℔ſſ. calidè, & ſudet horæ unius ſpatio.

His finitis, ſi puſtulæ adhuc non evanuerint, ſequenti Linimento partes affectæ inungantur.

℞ *Unguent. ex Oxylapatho* ℥ij. *Unguent. Pomati* ℥j. *flor. Sulphur.* ʒiij. *ol. Rhodii* ℈ſs. *f.* Linimentum.

Veruntamen Medicamenta præſcripta haud uſurpentur, niſi Venæſectione & Catharſi ritè præmiſſis.

## *De Angina.*

Angina inter Ver & Æſtatem ut plurimum aggreditur. Sequuntur febris inſultum dolor & faucium inflammatio, unde à tumore uvulæ, tonſillarum & laryngis, nec deglutire queat Æger, nec ſpiritum ducere.

Mittatur Sanguis è Brachio copiosè. Tangantur partes inflammatæ *melle Roſaceo, Sp. Vitrioli, vel Sp. Sulphuris ad ſummum acorem permixto, vel Sp. Vitrioli ſolo.* Mox Gargariſma ſequens uſurpetur, non vulgari modo; ſed ita ut ſine ulla agitatione in ore contineatur, donec incaleſcat; dein expuatur, & repetatur ſubindè.

℞ *Aq. Plantag. Roſ. Rubr. & Sperm. Ranar. an.* ℥iv. *album. ovor. in aquam agitando redact.*

*redact.* N° *iij. Sacchar. Chryſtal.* ℥*iij. M. f. Gargariſma.*

*Emulſio Refrigerans* in Pleuritide deſcripta indies ſumatur.

Sequenti manè, niſi febris & dolor inter deglutiendum remiſerint, Venæſectio repetatur, Catharſi in diem poſterum rejectâ, aliâs Catharticum lenitivum exhibeatur.

Manente adhuc affectu (quod oppidò rarum) repetatur adhuc Venæſectio, & Catharſis; Epiſpaſticum amplum ac forte applicetur cervici poſt primam Venæſectionem. Omni manè, niſi diebus purgationi datis Enema refrigerans & Emolliens injiciatur; Diætâ tenui utatur æger, & lecto per horas aliquot quotidie abſiſtat.

In his omnibus Febribus, quas intercurrentes appello, ut etiam in illa, quæ ſtationaria eſt, cautè obſervandum, quòd æger à lecto abſtineat magnam diei partem, ut juſculis hordeaceis, avenaceis, ac id genus aliis victitet, & cereviſiam tenuem lupulatam & hydrogalam pro potu ordinario bibat.

## *De Morbillis.*

Infantes plerumque impetunt. Primo die rigent horrentque, calent & frigent alternatim. Secundo conſummatam febrim accerſunt. Ægritudo vehemens, ſitis, inappetentia, lingua alba (non verò ſicca) tuſſicula, capitis

pitis & oculorum gravedo, & ſomnolentia perpetua, acriter premunt, humor è naribus & oculis jugiter ſtillat (quæ lachrymæ certiſſimum morbillorum ſignum.) Huc accedunt ſternutatio, tumor palpebrarum paulò ante eruptionem, vomitus, diarrhœa, cum dejectionibus colore viridi in dentientibus præſertim. Symptomata ingraveſcunt ad quartum uſque diem, quo quidem die (nonnunquam quinto) in fronte & facie maculæ parvæ, rubræ, pulicum morſibus perſimiles prodeunt, quæ numero & magnitudine auctæ racematim coaleſcunt, & faciem maculis rubris majuſculis interſtinguunt: Hæ maculæ è parvulis papulis, ſupra cutim paululum elevatis conflantur, quarum protuberantia digito levi, vix oculorum acie deprehendi poſſit.

[illegible] pectus ventremque, crura deinde tibiaſque propagantur maculæ; in his autem partibus latæ & rubræ, ſuperficiem cutis non ſuperantes conſpiciuntur. Eruptione morbillorum non perinde deliniuntur Symptomata ac in variolis. Vomitus poſt eruptionem nullus; ſed tuſſicula & febris cum ſpirandi difficultate intenduntur, in oculos defluxio, ſomnolentia & inappetentia perſeverant. Die ſexto aut circiter frons & facies aſpereſcunt, emorientibus puſtulis, ruptâque cuticulâ, in reliquo corpore maculæ latiſſimæ ſunt, & maximè rubeſcunt. Octavo circiter die in facie evaneſcunt, & vix in reliquo

quo corpore permanent. Nono nullæ prorſus ſunt; per faciem verò & artus quandoque & totum corpus in farinam quandam & ſquammulas faceſſunt, & tunc temporis febris, ſpirandi difficultas & tuſſicula augentur. In adultis à calido regimine liveſcunt primò, mox nigreſcunt exanthemata.

℞ *Decoct. Pectoral.* ℔ *iſs. Syrup. Violac. & Cap. Ven. an.* ℥ *iſs. M. f. Apozema*; Cap. ℥ *iij.* vel *iv.* ter vel quater in die.

℞ *Ol. Amygd. dulc.* ℥ *ij. Syr. Violar. & Cap. ven. an.* ℥ *j. Sacchar. Cand. albiſſ. q. ſ. M. f. Eclegma*, de quo ſæpius lambat, præcipuè quando tuſſis urget.

℞ *Aq. Ceraſor. nigr.* ℥ *iij. Syr. de Mecon.* ℥ *j. M. f. hauſtus* ſingulis noctibus ſumendus, à primo morbi inſultu, uſque dum convaluerit æger, doſin augendo vel imminuendo pro ratione ætatis.

Æger ſe contineat in lecto per biduum poſt primam eruptionem.

Si poſt Morbillorum diſceſſum, Febris, reſpirandi difficultas, & alia Symptomata Peripneumoniam mentientia ſupervenerint, ſanguis liberaliter extrahendus è brachio, ſemel, bis, vel ter, prout opus fuerit, convenienti ſpatio interpoſito inter ſingulas Venæſectiones; Atq; etiam continuentur *decoctum Pectorale ſupra de-*

*ſcriptum*, ut & *Lohoch*, vel *Ol. Amygd. dulc. per ſe.* Circa diem *xij.* ab invaſione Cathartico lenienti purgetur.

Diarrhœa, quæ morbillos excipit, Venæſectione ſanatur.

## *De Variolis.*

Diſtinctæ ſunt, vel confluentes: & in diſtinctis invadunt rigor & horror, calor intenſus, capitis & dorſi dolor vehemens, vomituritio, in adultis ingens in ſudores propenſio (& hinc minime confluxuras eſſe licet conjicere) dolor ſub ſcrobiculo cordis ſi manu prematur, ſtupor & ſomnolentia, nonnunquam & inſultus epileptici (in infantibus præſertim) quibus ſi, dentitione peractâ corripiantur, in procinctu eſſe variolas poſſis prædicere, ita ut ſi fortè inſultum epilepticum ſub veſperam patiantur, ſequenti Aurorâ in conſpectum ſe daturæ ſunt variolæ, mites ut plurimum & rariſſimè confluentes. Quarto incluſivè ab invaſione die, nonnunquam ſeriùs, rarò tamen ante illum erumpunt, quo tempore vel minuuntur Symptomata vel prorſus evaneſcunt. In eruptione autem ſubrubræ primò puſtulæ, tenuiſſimarum acicularum puncta æquantes, ſparſim ſe produnt per faciem, collum, pectus, & corpus univerſum; tunc in faucibus dolor, qui cum puſtulis ſurgentibus augetur. Circiter octavum à primo inſultu diem intervalla puſtularum priùs

priùs alba rubere jam incipiunt, in tumorem attolli, & dolore tenſivo affici; extenduntur palpebræ, & ut veſica inflata, oculos claudunt. Proximæ à facie intumeſcunt manus, digitíque diſtenduntur; puſtulæ faciei antea læves & rubræ, jam aſperæ (primum maturationis indicium) & ſubalbidæ apparent, ſuccum inſuper luteum, colore à favo non abludentem evomunt, faciei & manuum inflammatio jam ſumma in puſtularum interſtitiis colorem exhibet floridum, & roſarum Damaſcenarum æmulum, quem quidem colorem, quo mitiores ſunt variolæ, eo magis tum puſtulæ tum cutis interjecta ad amuſſim exprimunt: Quæ faciem occupant puſtulæ, aſperiores quotidie ac magis flavæ pro maturationis ratione; quæ manus & reliquum corpus, minùs aſperæ albæque magis conſpiciuntur indies. Die undecimo faciei tumor & inflammatio recedunt, puſtulæ maturæ & juſtam magnitudinem (viz. piſi grandiuſculi) adeptæ exareſcunt deciduntque. Decimo quarto vel quinto pereunt funditus; manuum autem puſtulæ pertinaciores, albæ adhuc & recentes die uno aut altero moras trahunt & tandem dirumpuntur; cæteræ autem, faciei nimirum & corporis deſquammantur, & in facie ſquammulis hiſce furfuraceis ſuccedunt foveæ. Per totum morbi decurſum alvus vel prorſus conſtipatur vel rariſſimè dejicit. Maxima eorum pars, qui ex hoc morbo pereunt, octavo die in diſcretis, undecimo

cimo autem in confluentibus intereunt: Quippe cùm in diſcreto hoc genere promoventur ſudores tum cardiacis tum regimine calidiore, prædicto die facies, quæ turgida & per interſtitia puſtulis interjecta inflammari debuerat, è contra flaccida reperitur & albeſcens, puſtulis nihilo minus rubentibus & elevatis, etiam poſt ægri mortem; ſudor, qui huc uſque manaverat, derepentè evaneſcit, æger interim phrenitide corripitur, anxietate, jactatione, ægritudine vehementibus, urinam parcè reddit ac frequenter, & tandem pauciſſimarum horarum ſpatio ad plures migrat.

In Confluentibus eadem Symptomata (ſed atrociora) febris ſcilicet, anxietas, ægritudo, vomituritio, &c. immanius ſæviunt; licet non ita promptè, ac in diſcretis, in ſudores diſſolvitur æger. Diarrhœa nonnunquam eruptionem præcedit, & ad diem unum aut alterum poſt illam protrahitur, quod in diſcretis vix contingit. Tertio die aut citiùs, rarò tardiùs, erumpunt; & quo citiùs, eo magis confluunt. Quandoque ad quartum vel quintum diem differtur eruptio ob atrox aliquod ſymptoma, v. g. acutiſſimum dolorem in lumbis, nephritici æmulum, in latere pleuritici, in artubus rheumatici, in ventriculo cum ingenti ægritudine & vomitu. Non perinde, ac in diſcretis, mox ab eruptione remittunt ſymptomata, ſed & febris & alia ad multos etiam dies poſt eruptionem puſtularum ægrum diſcruciant: Nunc

eryſipelatis ritu, nunc morbillorum erumpunt, eruptionis licèt tempore planè diſtinguantur. Creſcente morbo, non in molem aliquam ſpectabilem attolluntur; ſed in facie ſibi invicem implicitæ, tanquam veſicula rubra, vultum omnem contegunt, quem maturius elevant quàm diſcretæ: poſtmodum ad inſtar pelliculæ albæ vultui agglutinatæ cutis ſuperficiem non admodum ſuperant. Præterlapſo die octavo, pellicula alba quotidie paulatim magis exaſperatur, colore fuſco: Intenſus magis eſt dolor cutis, donec tandem in truculentiori morbo non niſi poſt diem viceſimum latioribus laminis deſquammatur. Hoc interim obſervatu dignum, quòd quanto puſtulæ matureſcentes propiùs ad colorem ſubfuſcum vergunt, eò deteriores lentiùs abſcedunt; quo magis autem flaveſcunt, eò minus confluxerint, & ocyùs ſe propipiunt. Ubi primùm deciderit pellicula, nullâ ſcabritie vultus afficitur; at verò mox illam excipiunt ſquammulæ furfuraceæ, indolis perquam corroſivæ, quæ foveas relinquunt patentiores, & ſæpe cicatrices. Quandoque humeri & dorſum cuticulâ denudantur. Morbi periculum à puſtularum frequentia & numero in facie ſolâ, æſtimandum. Puſtulæ manuum & pedum majores, & quo altiùs ab extremitate artuum aſcenderis, eo ſenſim minores & contractiores extant. In adultis ſalivatio, in infantibus diarrhœa (licèt non ita certò) confluentium comes. Ptyaliſmus nunc cum eruptione

ptione ſe prodit, nunc biduo triduóve præterito, materiam primò tenuem, die undecimo viſcidiorem ægerrimè excreat. Siticuloſus eſt æger, voce raucâ, ſomnolentiâ & ſtupore ingentibus corripitur, inter bibendum ſubinde tuſſit, & potus per nares revertitur. Tunc ceſſat ut plurimum ſalivatio, cui niſi faciei intumeſcentia (quæ licèt hoc ipſo die pro morbi genio aliquantiſper remitti debeat, non tamen niſi poſt diem unum alterúmve omnino recedere) & manuum jam inſigniter incipiens & diutiùs perdurans ſuppetias tulerint, ægro protinus moriendum eſt. Diarrhœa non ita maturè infantes occupat ac ptyaliſmus adultos. In utraque variolarum ſpecie febris ab inſultu ad eruptionem dominatur, inde dat inducias ad puſtularum maturationem, quâ peractâ, deſinit. Superveniunt à malo regimine ſymptomata anomala; qualia ſunt, puſtularum flaccidiras & ſubſidentia, phrenitis, coma, maculæ purpureæ puſtulis interſperſæ, & in earum ſummitate maculæ nigræ exiguæ in medio ſubſidentes, ſanguinis mictus & ſputum ſub initio morbi, urinæ ſuppreſſio.

Separationis tempus cum febrili ebullitione tribus vel quatuor primis diebus tranſigitur: Expulſio per reliquum morbi tempus, mediantibus parvis abſceſſibus, in carne perficitur.

Dies in quo maximè periclitatur æger, in confluentibus in minimè crudo & maximè vulgari genere, à primo morbi inſultu, eſt undecimus,

ecimus, in crudiore adhuc decimus quartus, in crudiſſimo decimus ſeptimus; nonnunquam tamen ſed rarius, non ante diem viceſimum primum moritur æger. Quinimo per patium illud omne, quod inter dies undecimum, & decimum ſeptimum interjacet, appetente veſperâ, moleſtiſſimo inquietudinis paoxyſmo, quotidie vexatur æger.

Mittatur ſanguis quovis ex tribus primis dibus ab invaſione ad ℥ *ix.* vel *x.* & poſtea *Vomitorium* exhibeatur *ad* ℥ *j.* vel ℥ *iſs. Infuſionis Croci Metallorum.*

His primis diebus diluatur ſanguis frequenti potatione Cereviſiæ tenuis lupulatæ. Cui Virioli Spiritus immiſceatur uſque ad plenam puſtularum eruptionem.

Cum variolæ omnes eruperint (quod ut plurimum eſt ſexto die poſt invaſionem) veſperi propinetur *Syr. de Mecon.* ℥ *j.* quæ repetatur dehinc ſingulis noctibus ad diem decimum poſt invaſionem. Eâ nocte, ſi Variolæ è Confluentium genere fuerint, augeatur doſis ad ℥ *iſs.* quæ dein repetatur ad ℥ *j.* manè, & ad ℥ *iſs.* ſerò ſingulis noctibus, donec æger convaluerit.

Si Syr. de Mecon. minus congruat, ſubſtitui poſſit *Laudan. Liquid.* viz. *gutt. xviij. pro* ℥ *j. Syrupi, & gutt. xxv. pro* ℥ *iſs.* Hoc interim obſervandum, quod ſi paregoricum bis in die exhibitum orgaſmum compeſcere non valeat (quod in variolis inſigniter confluentibus ſub finem

finem earum ſæpe accidit) tum 8ª quaque horâ exhibendum, vel ſæpius, ſi opus fuerit.

At ſi variolæ interſtinctæ fuerint, ſatis erit, ut ſolummodo ſingulis noctibus poſt plenam eruptionem exhibeatur Paregoricum, atque etiam in minori doſi.

Qualiſcunque autem generis variolæ fuerint, atque etiam qualicunque morbi tempore, ſi Phrenitis acciderit, omnino oportet ut inordinato ſpirituum motui frænum injiciatur, ac proinde ſi prior Paregorici doſis metam non attingat, alia atque alia aſſumatur, donec tumultus penitus compeſcatur, interpoſito inter ſingulas doſes ſpatio aliquo; ut pernoſcere valeamus, utrùm doſis ultima id, quod quærimus, præſtiterit, antequam nova ingeratur.

Si omnimoda urinæ ſuppreſſio acciderit, æger e lecto eximatur, & aliquantulum obambulet.

Si ſaliva ob calorem ſit adeò viſcida, ut expectorari non poſſit, tum Gargariſma cum ſyringa ſæpe in fauces injiciatur. Componatur *è Cereviſia tenui* vel *aq. hord. cum Melle Roſarum*; vel ſequens uſurpari poterit.

℞ *Cort. Ulmi* ℥vj. *rad. Glycyrrhiz.* ℥ſs. *Paſſul. enucleat.* № xx. *roſ. rub. p. ij. coq. ſ. q. aq. ad* ℔iſs. *in Colatur. diſſolv. oxymel. ſimp. & mell. Roſar. an.* ℥ij. *M. f. Gargariſma.*

Si epiſpaſtico opus fuerit, amplum ſatis & acre nuchæ applicetur nocte iſtâ, quæ magnam morbi criſin præcedit, mox ab aſſumpto paregorico;

gorico; plantis quoque pedum applicari poteſt allium, à die octavo, uſque ad morbi finem, ſingulis diebus renovandum.

Si Infans de Symptomatis, dentitionem comitantibus, jam ſecurus, derepente ſpaſmo corripitur, hic cogitandum eſt convulſionem iſtam, fortaſſis à naturæ conatu, quæ variolas, ſcarlatinam Febrem, aut morbillos, in corporis habitum protrudere ſatagit, pendere; applicetur itaque Emplaſt. Epiſpaſticum nuchæ, lecto quamprimum commıttatur æger, exhibeatur inſuper cardiacum, cui paregorici pauxillum eſt admiſcendum, puero nempe trienni *Laud. Liquid. gutt. v. in aq. Epidem. cochl. j.*

Si die undecimo, vel alio quovis poſt eum, febris ſecundaria cum inquietudine, jactatione, & cæteris id genus ſymptomatis, uſque adeò efferatur, ut Paregoricis utcunque repetitis ſuffiaminari nequeat, & mors quaſi immineat, tum ſanguis ad ſatis magnam quantitatem illicò detrahatur, nempe ad ℥*xij.* vel circiter, atque porrò repeti poſſit ſemel vel bis in ſequentibus diebus, ſi ſymptomatum dictorum ratio idem poſtulârit; at non aliàs. Catharſis etiam blanda die decimo tertio (non antea) vel quovis alio die eum inſequente (modo Venæſectio præceſſerit) inſtitui poſſit, nempe ex *Electuarii Lenitiv.* ℥*j. diſſolut. in aq. Cichor.* vel *lactis Alexeter.* ℥*iv.* Veruntamen nec Venæſectio nec Catharſis impediunt, quo minùs Paregorica in uſum veniant, quæ, (haud obſtantibus

ſtantibus iis) tam quoad largam doſin quàm repetitam exhiberi debent, ſi opus adhuc fuerit: nam ad hæc in hoc morbo ſemper confugiendum.

Quando Puſtulæ perfectè exaruerint, inungatur facies Linimento ex æqualib. partib. *ol. Amygd. dulc.* & *Pomat.* per biduum; at non ultra.

Die *xxj.* poſt invaſionem mittatur ſanguis è brachio; & ſequenti die Catharſis celebranda, quæ porrò alternis diebus ad tres alias vices repetatur.

Quoad Regimen; è lecto abſtineat ad ſextum diem poſt invaſionem, in quo poſtea ſe contineat uſque ad *xvij.* non aliis ſtragulis tectus, quàm quibus ſanus uti ſolebat.

Juſculis avenac. hordeac. Pomis coct. & Co reviſiâ tenui victitet, & poſt undecimum diem vini Hiſpanic. Cochl. *iv.* vel *v.* bis in die exhibeantur, ſi æger voluerit.

Si crurum tumor non evacuationibus præſcript. cedat, tum fotu fol. Malv. verbaſc. Sambuc. Lauri, cum floribus Chamæmel. & Melilot. in lacte coct. facilè fugatur.

Si primis morbi diebus æger Hæmoptoë laboret, vel urinam reddat ſanguineam, tum pulvis & tinctura, quæ in capite de Hæmoptoë præſcribuntur, ſextâ quâque horâ (donec ſymptomata iſta prorſus ceſſaverint) diligenter exhibeantur. Quo etiam caſu Paregorica liberali manu propinanda.

De

*De choreâ Sancti* Viti.

Convulſionis ſpecies quædam eſt, pueros puellaſve à decimo ætatis anno ad pubertatem invadit, prodit ſe claudicatione, vel inſtabilitate alterutrius cruris, quod æger poſt ſe trahit fatuorum more; manum ejuſdem lateris, vel pectori vel alii alicui parti applicitam, nequit continere in eodem ſitu horæ momento, ſed in alium ſitum & locum convulſione diſtorquebitur, quicquid æger contra nitatur. Antequam poculum ad os poſſit adducere, mille geſticulationes, circulatorum inſtar, exhibebit: Non enim rectâ lineâ ori admovet, ſed deductâ à ſpaſmo manu, huc illuc aliquandiu verſat, donec tandem fortè fortunà labris propius apponens liquorem derepentè in os injicit, & avidè haurit; deditâ tanquam operâ, ut ſpectantibus riſum moveret.

Sanguis ad ℥ *viij.* plus minus, pro ætate, ex brachio educatur.

Die ſequenti, dimidia pars, vel quiddam amplius, pro ratione ætatis, potionis communis hauriatur; & veſperi hauſtulus ſequens propinetur.

℞ *Aq. Ceraſor. nigr.* ℥ *j. aq. Epilept. Langii* ℥ *iij. Theriac Andromach.* ʒ *j. Laudan. Liquid. gut. viij. f. hauſtus.*

*Potio*

*Potio Cathartic.* ad tres vices, alternis diebus, repetatur; & *hauſtus Paregoricus* iiſdem noctibus.

Poſtea Sanguis iterum extrahatur; & ad Catharſin, uti priùs, æger revertatur, & ſic ſanguis mittatur, & alvus ſubducatur ad tertiam quartamve vicem; ſpatio tamen inter alternas evacuationes obſervato, ut nihil periculi ægro immineat.

Diebus à purgatione vacuis ſequentia præſcribantur.

℞ *Conſerv. Abſinth. Roman. & flaved. Aurantior. an.* ℥j. *Conſerv. Anthos* ℥ſs. *Theriac. Andromach. vet. & nuc. moſchat. condit. an.* ℥iij. *Zinzib. condit.* ℥j. *Syr. è ſucc. citr. q.ſ.f. Electuar.* de quo capiat ad magnitud. nuc. moſchat. manè & horâ quintâ pomeridianâ, ſuperbibend. vini ſequentis Cochlear. v.

℞ *Rad. Pœoniæ, Enul. Campan. Imperator. & Angelic. an.* ℥j. *fol. Rut. Salv. Betonic. Chamædr. Marrub. alb. & ſummitat. Centaur. min. an. Mj. Baccar. Juniper.* ℥vj. *Cortic. duor. Aurantior. Incidantur & infundantur frigidè in vini Canarini* ℔vj. Coletur tantummodo usûs tempore.

℞ *Aq. Rut.* ℥iv. *aq. Epilept. Lang. & Bryon. comp. an.* ℥j. *Syr. Pœon.* ℥vj. *M. f. Julap.*

cap.

cap. Cochl. iv. ſingulis noctibus cubitum iturus, inſtillando *Sp. C. C. gut. viij.*

Applicetur plantis pedum Emplaſt. è Caranna.

Ob Recidivæ metum pertundatur ei vena, & ad dies aliquot purgetur, eâ ipsâ anni ſequentis tempeſtate (vel paulo ante) quâ morbus invaſit.

Veriſimile mihi videtur, quòd hæc methodus curationi Epilepſiæ Adultorum convenire poſſit, quod tamen non adhuc expertus ſum. Cum verò Chorea *Viti* ætates teneras adoriri ſoleat, in Epilepſia Adultorum tam ſanguinis detrahendi quantitas, quàm Catharticorum doſis adaugeantur.

## *De Apoplexia.*

Somnus eſt profundiſſimus, & omnimoda ſensûs & motûs privatio, exceptâ reſpiratione, quam ægri habent difficilem cum ſtertore.

Mittatur Sanguis è brachio ad ℥ *xij.* quamprimum, ac poſtea ex venis juglaribus ad ℥ *viij.* deinde *Vomitorium ex* ℥ *iſs. vel* ℥ *ij. Infuſ. Croc. Metal.* propinetur ſtatim.

*Emplaſtr. Epiſpaſtic.* ſatis amplum ac forte cervici applicetur.

Dum hæc aguntur, erectus ſedeat æger in lecto, nec ſtragulis nimis obrutus.

*Sp. Sal. Armoniac. optimè rectificatus* admoveatur naribus.

Finitâ

Finitâ Vomitorii Operatione, ſubinde exhibeantur Cochl. iij. vel iv. ſequentis Julapii.

℞. *Aq. Rutæ* ℥iv. *aq. Bryon compoſ. & Antiepilept. Lang. an.* ℥j. *Sp. C. C. gut. xx. Sacchar. Cryſtal. q. ſ. f. Julap.*

Vel *Cochleare j. Sp. Lavendulæ compoſ.* per ſe propinari poſſit bis vel ter, durante paroxyſmo, horâ dimidiâ vel integrâ interpoſitâ.

Cavendum eſt, ne Cardiaca plus juſto calida, ac nimis frequenter (uti mos eſt) exhibeantur; nam quacunque demum ſpecificâ virtute pollere videantur, plus damni quam emolumenti ægro apportant, utpote quæ humores fundere (ac proinde malum augere) apta nata ſunt: quod etiam incommodum non minùs infertur ex nimia ſtragulorum congeſtione.

Paroxyſmo jam finito, ad præcavendam recidivam ſequentia exhibeantur.

℞ *Pil. Coch. maj.* ℈ij. cap. tertiâ quâque die per 6 vices horâ 4ª matutinâ, ſuperdormiendo.

℞ *Conſerv. flor. ſalv. Roriſmarin. an.* ℥ſs. *Conſerv. flaved. Aurant.* ℥vj. *nuc. moſchat. condit. & Zinzib. condit. an.* ℥ſs. *Theriac. Androm. vet.* ℥ij. *pulv. elect. diamb. diamoſch. dulc. an.* ℥j. *cum ſ. q. Syr. conditur. citr. f. opiata*, capiat ad magnitudinem Caſtaneæ manè & ſerò, ſuperbibend. *aq. Epilept. Lang. Coch.* ij.

℞ *Ambr.*

℞ *Ambr. griſ.* ℥ſs. *ol. deſtil. ſem. Aniſi, cinamom. nuc. moſchat. an. gutt. ij. ol. Caryophyl. ut. j. Sacchar. in aq. flor. aurantior. ſolut.* ℥iv. *f. s. a. Tabellæ*, quarum capiat unam ad libitum.

Ab uſu potûs generoſioris cujuſvis ſibi temperet, veſcatur juſculis avenaceis, hordeaceis, ut & ex pullis gallinaceis confectis, & quandoque etiam (præcipuè inter purgandum) carne pullorum, agnorum ac id genus ciborum cupeptωn.

## *De Ophthalmia.*

Ex oculorum inflammatione patet.

Mittatur Sanguis è brachio ad ℥x. Sequenti die ſumat potionem purgativam communem, quæ repetenda eſt per duas alias vices, duobus diebus interpoſitis; iiſdem noctibus ſumat *hauſtum paregoricum ex Syr. de Mecon.* ℥j.

Diebus à purgatione vacuis ſumat ter aut quater in die ℥iv. *Emulſionis ex ſem. frigid. maj. & Papav. alb.*

℞ *Aq. Plantag. Roſ. rubr. & Spermat. Ranar. an.* ℥j. *pulv. Tutiæ præparat.* ℥j. *f. Collyrium*, cujus guttæ aliquot inſtillentur oculis bis in die; at non incipendum, niſi poſt primam purgationem.

Si his remediis affectus haud cedat, repetatur Venæſectio ſemel vel bis (præcipuè ſi ſanguis

guis Pleuriticus apparuerit) ut etiam Catharsis.

Abstineat ab usu vini & cujusvis liquoris generosi, ab usu etiam ciborum coctu difficilium, & sale conditorum; & diebus inter purgationes bibat hydrogalam coctam.

Nota, quod sæpè accidit, quòd Venæsectioni & Catharsi utcunque repetitis Ophthalmia non cedit; hoc in casu *Paregoric. ex Syr. de Mecon.* ℥j. singulis noctibus exhibitum, curationem sine aliorum suppetiis absolvit.

## De Uteri Procidentia.

℞ *Cort. Querc.* ℥ij. *Coq. in Aq. font.* ℔iv. *ad* ℔ij. *sub finem addendo Cortic. Granator. contus.* ℥j. *ros. rubr. flor. Granator. an.* Mij. *dein adde vini rubr.* ℔ß. Colatura sit pro fotu, qui cum pannis laneis parti affectæ admoveatur manè horis duabus, antequam è lecto assurgat, & nocte post decubitum, donec symptoma penitus evanuerit.

## *De Paroxysmo Nephritico.*

Adsunt dolor fixus in lumborum regione, urina cruenta, arenularum aut calculi excretio, stupor cruris è directo renis laborantis, testis retractio ejus lateris, in quo est dolor, nausea & vomitus. Dolor colicus nephritico haud absimilis; licèt in quibusdam symptomatis planè

planè discrepet, quæ recensita in *Capite de Colica biliosa* videas.

Si Æger temperamento sanguineo fuerit, mittatur sanguis è brachio affecto reni respondenti ad ℥x. dein *liquoris possetici cui Rad. Althææ* ℥ij. *incoctæ fuerint Cong. j.* quam citissimè deglutiatur; & Enema sequens injiciatur.

℞ *Rad. Althææ & Lilior. an.* ℥j. *fol. Malv. Parietar. Branc. Ursin. & flor. Chamæmel. an. M. j. Sem. Lini & Fœnugr. an.* ℥ss. *coq. s. q. aq. ad* ℔iss. *M. F. Enema.* Rejectis Vomitorio & Enemate, *Laudani Liquidi satis magna dosis* propinetur, viz. *ad gutt. xxv.* vel *Pill. Matth.* gr. *xv.* vel *xvi.* In senibus verò, & morbo diuturno fractis, ut & vetulis Vaporibus obnoxiis (præcipuè si sub initium Paroxysmi urinam eæ reddant nigram & sabulosam) Venæsectio omittatur; at quoad reliqua, eâdem quæ dicta est viâ omnino procedendum.

*In mictu Sanguineo à calculo renibus impacto.*

Capiat æger, statò quodam die, singulis hebdomadis, *Mannæ* ℥iiss. in seri lactis ℔ij. solut. Conducit aliquando, haustus cerevisiæ tenuis liberalior.

In calculo renum grandiori, dolor hebes, & obtusus, nullatenus auctus, sed satis tolerabilis persentitur, nec paroxysmus nephriticus comitatur.

Aquis chalybeatis æger ſe non tutò committet, niſi prius certò ſciat, calculum majorem non eſſe, quam ut per ureteres viam poſſit invenire, quod, hoc tantum modo, certò poteſt dignoſci, ſi nempe, jam antea, paroxyſmum aliquem nephriticum fuerit perpeſſus, (cum dolore ſcilicet ſæviſſimo in renum alterutro, per ureterum ductus propagato, & vomitu enormi) pro comperto habeat, non tam grandiorem aliquem calculum pelvi concludi, quam lapillorum exiguorum congeriem, quorum unus ſubindè in ureterem illabitur, atque ita paroxyſmum invitat, qui fere non deſinit, niſi calculo in veſicam detruſo; cum ſic ſe res habeat, nullum eſt magis efficax remedium, quam aquarum chalybeatarum potatio: quod ſi calculoſus, nullo unquam paroxyſmo nephritico fuerit tentatus, cum jam major ſit calculus, quam qui ex pelvi poſſit elabi, devitandæ ſunt aquæ ferreæ.

## *De Dyſenteria, Diarrhœa & Teneſmo.*

Rigent horrentque, ſequitur deinde totius corporis calor, quem brevi excipiunt ventris tormina, & hæc, dejectiones crebræ, & mucoſæ omnes (niſi quòd ſtercoroſæ nonnunquam interponantur) cum ingenti cruciatu & moleſtiſſimo viſcerum omnium quaſi deſcenſu, quoties exoneratur alvus, ſanguinis interim lineamentis intertextæ, licèt aliquando ne minimum

mum ſanguinis per totum morbi decurſum admiſceatur. Progreſſu morbi ſanguis ſincerus quandoque excernitur, & gangrænâ inſanabili afficiuntur inteſtina. Si æger ætate floreat, vel cardiacis incaleſcat, ſebricitat admodum, lingua eſt ſubalbida, quadam mucilagine densè obſita, interdum nigra & ſicca. Proſternuntur vires, diſſipantur ſpiritus, aphthæ inſuper oris interna fauceſque obſident, præſertim ubi materiæ peccantis evacuatio adſtringentibus præpoſtere impeditur, non exacto per cathartica morbi fomite. Aliquando, exulante febre, agmen ducunt tormina, & cætera morbi ſymptomata ſequuntur.

In diarrhœa humores excrementitii ſine ſanguine aut inteſtinorum exulceratione excernuntur.

In teneſmo aſſidua eſt deſidendi cupiditas, nihil tamen præter mucoſum quoddam & ſubcruentum aut purulentum rejicitur.

Mittatur ſanguis è brachio quamprimum, eademque nocte *Paregoricum* propinetur; & mane ſequenti *Potio purgans communis*, repetenda alternis diebus ad duas alias vices, & *Paregoric.* poſt Purgationes, quam primum ſuo defunctæ ſunt munere; ac diebus à purgatione vacuis idem manè & ſerò eſt exhibendùm.

Poſt Venæſectionem & Catharſin ſemel celebratam, per totum morbi decurſum, ſequens Cardiacum exhibeatur.

℞ *Aq. Ceraſ. nigr. & Fragor. an.* ℥iij. *Aq. Epidem. Scord. comp. & Cinam. hordeat. an.* ℥i. *Margarit. præparat.* ℥iſs. *Sacchar. cryſtal. q. ſ. adde Aq. Roſar. Damaſcen.* ℥ſs. *ad gratum ſaporem, m. f. Julap.* de quo cap. cochl. *iv.* vel *v.* in languoribus, vel ad libitum.

Potus ſit Lac coctum cum *Aq. Triplo*, vel *Decoct. Alb.* ut,

℞ *C. C. C. & mic. panis alb. an* ℥ij. *coqu. in. Aq. font.* ℔iij. *ad* ℔ij. *poſtea ſ. q. Sacchar. Albiſſ. edulcoretur.* Vel ſi dejectio virium poſtulaverit, *Aq. font.* ℔ij. *cum Vin. Canarin.* ℔ſs *ſimul cocta*, pro potu frigidè aſſumantur.

Cum per tres vices purgatus fuerit æger, tota curatio uſui *Laudani* bis vel ter in die repetiti poſtea incumbit, niſi quòd *Enema ex Lac. vaccin.* ℔ſs. *& Theriac. Androm.* ℥iſs. ſubinde injiciatur, quod in alvi egeſtionibus mire confert.

Cum alvi fluxus in Diarrhœam tantum aſſurgat, Phlebotom. & Catharſi præſcript. omiſſis, ſequ. Bolus ſingulis matutinis exhibeatur.

℞ *Rhabarb. pulv.* ℥ſs. (*plus minus pro ratione virium*) *cum ſ. q. Diaſcordii f. Bolus, addendo Ol. Chym. cinnam. gutt. ij.*

Noctibus inſequentibus *Paregoric. ex Aq. cinnam. hordeat* ℥j. *& Laudan. Liquid. gutt. xiv. exhibeatur.*

Ubi tormina ſicca ſine dejectionibus excruciant, curantur diluendo ſero lactis frigide pro-

propinato, & tepidè injecto; ut in Cholera morbo, pullorum juſculo aut lacte cereviſiato.

Si hic morbus diutius excurrat, adeò ut omnis ejus vis in rectum inteſtinum decumbat, cum aſſidua deſidendi cupiditate, Diæta Analeptica inſtituatur, & liquor aliquis Cardiacus pro libitu ingeratur, ut reſtaurari poſſint vires, quibus redeuntibus, Teneſmus ſponte ſe proripit.

In dyſenteria malè curata dolores aliquando ægrum ad annos aliquot diſcruciant, in hoc caſu Phlebotomia repetita priſtinam ſanitatem reſtituet.

Notandum eſt, quòd in conſtitutionibus huic morbo minùs faventib. omiſſis evacuationibus dyſenteria Laudano ſolo ſanatur, quod manè & ſerò repetendum, donec ſymtomata penitus evanuerint, vel etiam ter Nycthemeri ſpatio, ſi opus fuerit.

## *De Colica bilioſa.*

Inteſtinorum dolor atrociſſimus eſt, per horas aliquot affligit febris in hunc morbum deſinens, qui aut quaſi injectâ faſciâ conſtringit aut in punctum contractus quaſi terebello perforat; ſubinde remittitur, mox recrudeſcit. Sub initio non ita certò ad unum aliquod punctum determinatur, neque tam crebra vomituritio, neque alvus ita pertinaciter catharticorum vim eludit; quo autem magis auge-

tur dolor, eo obſtinatiùs in puncto figitur, vomituritio frequentior & major alvi adſtrictio, colicâ tandem in iliacam tranſeunte. A nephritide ſic diſtinguitur. 1. Dolor nephriticus fixus in rene eſt, & ab eo ad teſtem exporrigitur ſecundum longitudinem ureteris. Colicus vero vagus eſt, & cinguli inſtar medium ventrem affligit. 2. A paſtu colicus dolor augetur: Nephriticus potius imminuitur. 3. In colico dolore, ab excretione per alvum & per vomitum majus levamen emergit, quàm in nephritico. 4. In dolore nephritico urina primùm clara & tenuis, poſtea aliquid ſubſidet, & tandem arenulæ aut lapilli excernuntur: In colica urinæ ab initio craſſiores exiſtunt.

Sanguis è brachio paulò liberaliùs educatur, & poſt horas *iij.* vel *iv. Anodyn.* exhibeatur. Poſtero die *Catharticum Lenitiv.* & interpoſito die uno, ad tertiam vicem propinetur.

At ſi malum hoc fructibus horariis, vel aliis cibis avidè ingeſtis, originem debeat, tum ventriculus ante omnia lacte cereviſiato affatim hauſto eluendus eſt; quo exantlato Anodynum propinandum eſt; ſequenti autem die vena ſecanda eſt, & in ordine præſcripto procedendum.

Ubi hic morbus minùs peritè tractatus diutiſſimè fatigaverit, & ægrum penè confecerit, liberalior ſive aquæ Epidem. ſive Mirab. ſive aliûs cujuſcunque, qua ſanus maximè delectabatur, uſus, ægrum hoc tempore ultra ſpem omnem juvit.

De

*De cholera Morbo.*

Intra Auguſti limites ſe continens, vix in priores Septembris hebdomadas evagatur. A crapula & ingluvie excitatus ſimilis affectus, nullo temporis diſcrimine inſurgit, qui licèt eodem curetur modo, alterius tamen eſt ſubſellii. Adſunt vomitus enormes, ac pravorum humorum cum maxima difficultate & anguſtia per alvum dejectio; ventris ac inteſtinorum dolor vehemens, inflatio & diſtentio; cardialgia, ſitis; pulſus celer ac frequens, parvus & inæqualis; æſtus & anxietas, nauſea moleſtiſſima, ſudor, crurum & brachiorum contractura, animi deliquium, partium extremarum frigiditas, & ſimilia, quæ ægrum in 24 horarum ſpatio interimunt.

Pullus tenerior in tribus circiter Aq. font. Congii elixetur, ut carnis ſaporem vix perceptibilem liquor referat, hujus decocti vel (defectu ejus) liquoris poſſetici capaciores aliquot cyathos æger tepidè exhauriat; & eodem tempore ex eodem liquore *Enemata* ſucceſſivè repetenda. Hauſtibus & Clyſteribus *Syr. Lactuc. Violar. Portulac. Nymph. eorúmve*, alicujus uncia ſubinde admiſceri poterit.

Exantlato hoc Eluvionis penſo (quod *iij.* vel *iv.* horas ſibi vindicat) *Paregoricum* curationi Coronidem imponit.

Si non acceſſerit medicus, niſi poſtquam æger vomitu & dejectionibus per multas horas continuatis fuerit exhauſtus, & jam frigeſcant membrorum extrema, tum ad *Laudanum Liquidum* confugiendum eſt in paulò majori doſi (viz. *Laudan. liquid. gutt. xxv. in Aq. cinnam. fort.* ℥*j.*) & cum ceſſaverint ſymptomata, nihilominus mane & ſerò quotidiè eſt repetendum, at in minori doſi, donec priſtinas vires receperit.

Eſt ſpecies quædam choleræ morbi infantibus valdè familiaris, ac ſæpenumerò funeſta, ex Dentitione vel Crapula originem ducens.

Horum ætas tenella minime fert, ut diluatur ventriculus largâ liquorum potatione, multo minùs ut Catharticis tumultus major in humoribus excitetur, ſed Curationem *Laudano* ſolo aggredi oportet; quamobrem *gutt. ij. iij.* vel *iv.* vel *plures*, pro ratione ætatis, Cochlear. *j.* cereviſiæ tenuis vel Aquæ cujuſvis appropriatæ inſtillatæ exhibeantur, repetanturque, ut opus fuerit.

### *De Colica Pictonum.*

Eſt ſpecies quædam Colicæ, quæ degenerare ſolet in Paralyſin, & motûs omnimodam manuum ac pedum depravationem; quam hoc nomine *Riverius* in capite de Paralyſi indigitat, & apud inſulas Caribum notiſſima eſt; utpote quæ magnam vim hominum infeſtat.

Hunc

Hunc dolorem atrociſſimum ſanat *Balſam. Peruv.* frequenter ac in magna dôſi exhibitum, nempe ſi *ejus gutt. xx, xxx,* vel *xl. Sacchar. albiſſ. cochl. j. inſtillentur,* & bis vel ter in die dentur: at Paralyſis huic remedio haud cedit.

## *De iliaca Paſſione.*

Periſtalticus inteſtinorum motus invertitur, propinata cathartica & injecta enemata in emetica ſtatim tranſeunt, alvi fæces per vomitum rejiciuntur.

Mittatur ſanguis è brachio dextro ad ℥ *ix,* vel *x.* poſt aliquot horas exhibeatur ſequens Pulvis.

℞ *Reſin. Scammon. vel (defectu ejus) Reſin. Jalap. gr. xij. Calomelan. Torquet.* ℈*j. m. f. Pulvis*; cap è Cochl. *j.* lactis vaccini, ſuperbibendo ejuſdem lactis Cochl. *j.* vel alterum: vel ſi forma Pilularum magis arrideat:

℞ *Pil. ex duob.* ℥*ſs. Calomel. Torquet.* ℈*j. cum ſ. q. Balſam. Peruv. f. Pill. iv.* ſumend. c. Cochl. *j.* Syr. Violarum.

Sin autem æger hoc remedium per os rejecerit, *Laudan. Liquid. gutt. xxv. Aq. Cinnam. fort.* ℥*ſs.* inſtillatæ ſtatim propinentur; cum vero horum uſu vomituritio, ac ventris dolores, remiſerint, catharticum dictum iterum exhibeatur: verum ſi anodynorum virtu-

te ac vi jam penitus infractis vomiturition ac dolor iterum redierint, cathartico interim in corpore reſtitante, ad uſum paregorici ſupra deſcripti revertendum, idemque eò uſque repetendum quartâ vel ſextâ quâque horâ donec inteſtinis quies omnino inducatur, quo quidem tempore catharticum operationem ſuam per ſedes exeret.

Nec ſanè abs re fuerit elapsâ horâ unâ, aut alterâ, poſt Venæſectionem, clyſterem aliquem, ex iis qui potentiùs operantur, injicere, fumum ſcilicet nicotianæ, ex tubulo inverſo, per veſicam majuſculam, in inteſtina validiſſimè inſufflatum, quæ quidem operatio poſt aliquot horas reiterari debet, niſi citius meatus deorſum pateat.

Finitâ operatione Cathartici, ſumat ſequentem hauſtum.

℞ *Aq. Cinnam. fort* ℥j. *Laudan. Liquid. gutt. xxv. m. f. Hauſtus,* qui porrò repetendus eſt bis, ter, vel ſæpius in die, donec vomituritio & dolor penitus ceſſaverint; atque etiam, iſtis ſymptomatis ceſſantibus, nihilominus ad curationem confirmandum per aliquot adhuc noctes capiat idem paregoricum in minori doſi cubitum iturus.

### *De Fluxu Menſium immodico.*

Menſtruorum fluxus, naturaliter conſtitutus, capacitatem teſtæ ovi anſerini implet.

Patet

Patet autem immodicus ex difficili tolerantia, virium lapſu, inappetentia, cachexia, colore cadaveroſo, & pedum intumeſcentia.

Mittatur ſanguis è brachio ad ℥ *viij.* ſequenti manè ſumatur *potio purgativa communis*, repetaturque alternis diebus ad duas alias vices, & noctibus poſt purgationem *Paregoric. è Syr. de Mecon.* exhibeatur.

Diebus à purgatione vacuis ſumat ſequentia.

℞ *Conſerv. Roſar. ſicc.* ℥ *ij. Trochiſc. de terr. Lemnia* ℥ *iſs. Cort. Granator. Corall. rub. præparat. & boli Armen. an.* ℥ *ij lapid. hæmatit. Sanguin. Draçon. an.* ℥ *j. cum ſ. q. Syr. Cydonior. f. Elect.* de quo ſumat ad magnitudinem nucis moſchat. maj. manè & horâ quintâ pomeridianâ, ſuperbibendo ſequentis Julapii Cochl. *vj.*

℞ *Aq. Germin. Querc. & Plantag. an.* ℥ *iij. Aq. Cinnamom. hordeat; & Syr. de Roſis ſiccis an.* ℥ *j. Sp. Vitriol. q. ſ.* ad gratam aciditatem.

℞ *Fol. Plantag. & Urtic. aa q. ſ. Contundantur ſimul, & ſuccus exprimatur, & poſtea clarificetur*, de quo cap. Cochl. *iv.* vel *v.* ſæpiuſculè ad libitum.

Poſt primam purgationem applicetur emplaſt ſeq. regioni Lumborum.

℞ *Emplaſt*

℞ *Emplaſt. de Minio & ad Herniam an. part. æq. extendantur ſuper alutam, f. Emplaſtrum* regioni Lumborum admovendum.

## *De Colica hyſterica.*

Eſt ſpecies quædam, ſeu potiùs inſigne ac valdè uſitatum affectionis hyſtericæ ſymptoma, cui comitem ſe adjungit dolor circa ſcrobiculum Cordis atrociſſimus, uti etiam humorum viridi colore per vomitionem ejectio.

Primo, quò ſaburra vitioſorum humorum in ventriculo contentorum minùs impediat Paregorici efficaciam, ſatis magna quantitas *liquoris poſſetici* affatim ingurgitetur, atque iterum per vomitionem reiciatur, nempe *ejuſdem liquoris Cong. iſs,* quo peracto *Laudan. Liquid. gutt. xxv. in Aq. cinnam. fort. Aq. Epidem. vel alius vehiculi* ℥*j.* propinentur.

In uſu Paregoric. hoc notandum, quòd eò uſque repetendum eſt, donec ſymptomata omnia penitus ceſſaverint, debitum interea tempus interjiciendo; ut dijudicare valeamus, quid prior doſis effecit, antequam novam ingeramus.

Si ſanguineo temperamento præditus ſit æger, vel ſi antea hoc ſymptomate haud multum infeſtatus fuerit ac viribus fractus, hoc in caſu ſanguis primùm è brachio detrahendus antequam vomitorium præſcriptum propinetur.

Hâc

Hâc ipsâ Methodo Clavus hyſtericus curatur. Sin verò Colica Hyſterica diuturnior fuerit per Paroxyſmos affligens: tum in eorum intervallis, ſive extra Paroxyſmum, ſequ. uſurpentur.

℞ *Zedoar. ſubtiliſſ. pulv.* ℥j. *cum ſ. q. Syr. è conditur. citri f. Bolus*, ſumendus manè & ſerò per *xxx.* dies, ſuperbibendo ſequentem infuſionem.

℞ *Zedoar. inciſ. minutim* ℥ſs. *vini Canar.* ℥iv. *infundantur frigidè per xij horas; deinde colentur pro uſu*; vel *Balſam. Peruv.* ut in Colica Pictonum, uſurpetur.

Quæ de Colica Hyſterica hic dicuntur, de Colica etiam hypochondriaca dicta ſunto; ut & hoc diligenter advertendum.

Utriuſque affectûs natura ſibi vendicare videtur, ut alia atque alia tententur, donec tandem ad illud remedium perventum fuerit, cui quaſi genuinè morbus cedit; tametſi utplurimum Chalybs imprimis confert.

Colica Hyſterica tam in hypochondriacis quam hyſtericis ſæpiſſime in Iéterum deſinit, qui ſenſim ſponte ſuâ faceſſit, ſin autem longas moras nectat, exhibeatur Apozema ſequens.

℞ *Rad. Rub. Tinctor. curcum. an.* ℥j. *chelidon. maj. cum toto & ſummitat. centaur. m n. an.*

*an. M. j. coq. in æqualib. partibus Vin. Rhenani & Aq. fontan. ad* ℔ij. *colatur. diſſolv. Syr. de 5 radicibus* ℥ij. *m. f. Apozema.* de quo capiat ℔ſs. calidè manè & ſerò, uſque dum convalucrit.

*De Ictero, colicâ neutiquam prægreſſâ.*

Color flavus in toto corpore, præſertim in oculorum albis eſt conſpicuis, quo quidem colore objecta omnia apparent affectus tincta, corporis pruritus, ſegnities, & laſſitudo, linguæ amaror, vomitus bilioſus interdum, ſingultus, egeſtio alba, urina crocea & pannum immerſum eodem colore inficiens, morbum indigitant.

Exhibeatur *potio purgativa communis,* dein aggrediatur uſum ſequentium; & durante eorundem uſu, Cathartieum præſcriptum quarto quoque die exhibeatur.

℞ *Conſerv. Abſinth. Rom. conſerv. flaved. Aurant. an.* ℥j. *Angelic. condit. nuc. moſchat. condit. pulv. Ari compoſ. & chalyb. cum aceto præparat. an.* ℥ſs. *Extract. centaur. min. Extract. Gentian. & cremor. Tartar. an.* ℥ij. *croc. Anglic. pulv.* ℥ſs. *cum ſ. q. Syr. de 5 radicibus f. Electuarium,* de quo cap. ad magnitud. nuc. moſchat. manè & horâ quinta pomeridianâ (vel vice Electuarii ſumat pilulas chalybeatas) ſuperbibendo poſt matutinam doſin *aquarum mine-*

*mineralium purgantium* ℔iv. & poſt pomeridianam doſin *Apozemat. in colica hyſterica deſcript.* ℔ſſ.

At ſi ſigna hydropis affuerint, tàm manè quàm ſerò apozema ſuperbibat.

Si vel his diu continuatis morbus adhuc obſtiterit, vel *aquas ferreas* eundum eſt, quæ ex ipſo fonte ſunt bibendæ.

### *Ad præcavendum abortum.*

Eadem præſcribenda, ac in Fluxu menſium immodico, omiſſis ſolummodò catharſi & ſuccis.

### *De immodico Lochiorum fluxu.*

Fluxum lochiorum naturalem tria conſtituunt, ſanguis ſcilicet purus, copioſus, per triduum primum fluens; dehinc ſanguis aquoſus, loturæ carnium craſſiori ſimilis, per quatriduum præterpropter durans; hinc poſtmodum materia viſcoſa & mucilaginoſa, pauco aut nullo ſanguine permiſta, per quinque aut ſeptem dies prodiens, & ſic porro ad menſis finem.

Fluxus autem eorum immodicus dijudicandus eſt ex virium dejectione, animi deliquio, ſanguine grumoſo effluente, cibi faſtidio, dolore in hypochondriis, ventris tenſione, pulſu debili

debili & frequenti, visûs obtenebratione, aurium tinnitu, & convulsionibus.

Victus incrassans instituatur, & potus sequens addatur,

℞ *Aq. Plantag. vim. rubr. an.* ℔j. *coq. simul ad tertias, & deinde edulceretur ſ. q. ſaccha. albiſſ.* cujus, cum frigeſcat, cap. ℥ſs. bis vel ter in die.

Interim Julap. aliquod Hyſtericum (at ſatis mite) ſubinde propinetur, & nodulus ſequens naribus applicetur.

℞ *Galban. Aſa fœtid. an.* ʒij. *caſtor.* ʒiſs. *ſal. ſuccin. volatil.* ʒſs. *m. f. nodulus,* vel

℞ *Sp. ſalis Ammoniac.* ʒij. quem ſæpe olfaciat.

*De ſuppreſſione Lochiorum.*

Intumeſcit venter, dolor gravativus imum ventrem, lumbos & inguina occupat, facies rubeſcit, difficilis eſt reſpiratio, perturbantur oculi, rigores, febres acutæ adſunt, animi deliquia, ſudores frigidi, pulſationis & ardoris ſenſus in utero, paralyſis in partibus inferioribus, ut & interdum epilepſia.

E veſtigio lecto addicatur, & *Emplaſt. hyſtericum* umbilico admoveatur quamprimum. Elect. ſeq. adhibeatur.

℞ *Con-*

℞ *Conſerv. Abſinth. Rom. Rut. an.* ℥j. *Trochiſc. è Myrrh.* ℥ij. *Caſtor. croc. Anglic. ſal. volatil. Sal. Armoniac. & Aſæ fœtid. an.* ℥ſs. *cum ſ. q. Syr. è 5 radicib. f. Elect.* cap. ad magnitud. nuc. moſchat. quartâ quâque horâ, ſuperbibendo ſeq. Julap. Cochl. 4 vel 5.

℞ *Aq. Rutæ* ℥iv. *Bryon. compoſ.* ℥ij. *Sacchar. cand. q. ſ. f. Julap.* vel ſumat *Trochiſc. de Myrrh.* ℈j. quartâ quâque horâ.

Si tota prædictorum quantitas conſumatur, & Lochia adhuc ſupprimantur, *Laudanum* unicâ ſaltem vice exhibendum; ut, *Laudan. Liquid. gutt. xiv. in Aq. Bryon. compoſ. vel Laudan. ſolid. gr. iſs. cum Aſæ fœtid.* ℈ſs. *in pilulas duas redactum.*

Si hæc nihil proficiant, tum Enema ex lacte Saccharato, unicâ ſaltem vice, injiciatur.

## *De Hydrope.*

Foveæ à digitorum impreſſione in ſurarum parte inferiori ſub noctem conſpicuæ, manè evaneſcentes, hoc malum primùm indicant, maximè ſi ſpiritus interim ægriùs ducatur (haud rarò tamen fœminæ prægnantes, & quibus catamenia ſiſtuntur, ut & viri ab aſthmate inveterato derepentè liberati eodem tumore tentantur) pedibus cruribúſque ad extremum extenſis, in abdomen ruunt aquæ, & ad ultimum capacitatis ſuæ limen paulatim diſtendunt,

dunt, donec tandem in viſcera nobiliora impetu ab iis facto æger ſubmergitur. Pro ratione qua partes à morbo obſeſſæ mole augentur, aliæ magis magíſque gracileſcunt, urgentibus dypſnœâ, urinæ paucitate, & ſiti intensâ. Viros, ingraveſcente ætate; fœminas, ubi jam parere deſierint, utplurimum adoritur.

Indicationes curativæ dirigendæ ſunt primò ad aquarum evacuationem, deindè ad inſtaurandum ſanguinis robur; quò novus aquarum proventus præcaveri poſſit.

℞ *Vini albi* ℥iv. *Jalap. ſubtiliſſ. pulv.* ℥ſ. *Zinzib. pulv.* ℈ſs. *Syr. de Spin. Cervin.* ℥j. *M. f. Potio*, ſumenda ſummo manè, repetendaque ſingulis vel alternis diebus, pro ratione virium, donec partes affectæ detumuerint; vel,

℞ *Tamarind.* ℥ſs. *fol. Senn.* ℥ij. *Rhei* ℥iſs. *Coq. in ſ. q. aq. ad* ℥iij. *in Colatur. diſſolv. Mannæ & Syr. Roſ. ſolut. an.* ℥j. *Syr. de Spin. cervin.* ℥ſs. *Elect. è ſucc. Roſar.* ℥iij. *M. f. Potio*, ſumenda, ut ſupra; vel,

℞ *Pil. ex duob.* ℈j. *Elater. gr. iij. Ol. Chym. Caryophyl. gut. ij. f. Pil.* N° *iij.* cap. ſummo manè, repetanturque, ut opus fuerit; vel,

℞ *Gambogii gr. xv. vin. alb. & aq. Cichor. an.* ℥iſs. *Syr. de Spin. Cervin.* ℥ſs. *m. f. Potio* ſumenda ut ſupra; vel,

℞ *Cortic.*

℞ *Cortic. interior. Sambuc. è ligno deraſi M. iij. coq. in aq. & lact. partibus aq. miſtorum* ℔*ij. ad* ℔*j.* Colaturæ capiat dimidiam partem manè, & alteram ſerò quotidie, donec convaleſcat. At hoc remedium, niſi in corporibus, quæ facillimè purgativis cedunt, nihil confert.

Circa uſum Catharticorum in hoc morbo hæc tria advertenda ſunt.

Primo, utrùm ægri corpus facilè, an verò difficulter Catharticis cedere ſoleat, cùm in iis qui facilè purgantur *Syrup. de Spin. Cervin. ſolus ad* ℥*j.* exhibitus aquas ſat copiosè eliciat; in illis verò, qui difficulter purgantur, Cathartica ſupra deſcripta, utut valida, vix ſcopum attingant.

Secundò, Cathartica omnia, quæ ſegniùs operantur, magis officiunt quam proſunt; quamobrem purgatio paulò fortior & nimia, juſto debiliori eſt præferenda.

Tertio, Aquæ eâ maximâ celeritate exhauriri debent quam ægri vires ferre queant: ne nimis longa inter purgationes intervalla adhibendo, aq. denuò affatim colligantur.

Eſt autem ubi medicamenta omnia, qualiſcunque demum generis, quæ alvum movent, prorſus omitti poſſint ac debeant; quoties enim æger conſtitutione eſt infirmiori, vel fœmina vaporibus obnoxia, aquarum evacuatio Diureticis ſolùm committenda, inter quæ ea ſunt efficaciſſima, quæ ex ſalibus Lixivialibus fiunt; ut, ℞ *Ci-*

℞ *Cinerum Geniſtæ* ℔j. *infundantur frigidè in vini Rhenan.* ℔iv. *liquoris per filtrationem colati* ʒiij. manè, quintâ pomeridianâ & ſeptimâ ſingulis diebus exhibeantur, uſque dum humor evanuerit.

Aquis jam penitùs evacuatis Calefacientia & corroborantia medicamenta uſurpanda; ut,

℞ *Rad. Raphan. Ruſtican. fol. Cochl. hortenſ. Abſinth. vulgar. Salv. ſummitat. Centaur. min. & Geniſtæ, infundantur in cereviſia forti* pro potu ordinario, qui potus incipienti hydropi aliquando ſine Catharticis ſubvenit; vel ſequent. utatur;

℞ *Conſerv. Cochl. hortenſ. Abſinth. Roman. an.* ʒj. *Extract. Gentian. Abſinth. vulgar. & Centaur. min. an.* ʒiij. *cum ſ. q. Syr. è cortic. citr. f. Electuarium,* de quo ſumat ad quantitatem nuc. moſchat. maj. manè, quintâ pomeridianâ, & veſperi, ſuperbibend. ʒiv. ſequentis infuſionis.

℞ *Rad. Gentian.* ʒj. *ſummitat. Geniſt. Centaur. min. & Abſinth. vulgar. an. M. j. Sem. Fœnicul. & Petroſelini an.* ʒij. *inciſis minutim affunde vini Rhenani* ℔iv. *ſient ſimul in infuſione frigida,* & coletur ſolummodo tempore uſûs.

Notandum, cùm ad corroborantium uſum deventum fuerit, ægrum nullo modo eſſe purgandum,

andum, tantiſper dum ſanguinis corroborati-
ni navetur opera.

Quod & per ſalium Lixivialium uſum om-
ino obſervandum, quo quidem tempore pa-
iter corroborare convenit.

℞ *Rad. Raphan. Ruſtican. ℥ iij. fol. Cochl. Hor-
enſ. Abſinth. vulg. Salviæ, ſummitat. Centaur. &
Geniſtæ, an. M. iij. Aurant. inciſ.* N° *iij. Infun-
antur in Congiis vj. Cereviſiæ fortis non lupula-
æ, inter fermentandum*; ſit pro potu ordinario.

## *De Gonorrhœa virulenta.*

Corripiunt dolor genitalium inſolitus, & ro-
atio teſticulorum, in non recutitis macula
morbilli æmula glandem occupat, ſtatim à cu-
us exortu liquor, ſeminis inſtar, exſtillat, qui
uotidie à ſeminis colore & conſiſtentia rece-
ens flaveſcit tandem dilutius ovi vitello; in
ue virulentiore vireſcit, & aqueo humori per-
miſcetur ſanguine tincto. Puſtula dicta ulcus
it, aphthis infantum ſimile, indies latius &
rofundiùs depaſcens, ſuperveniente orificii
allo. Verpi non glandis ulcere ſolâ Gonor-
hœâ tentantur. Mox magnus dolor in pene,
uoties erigitur; quaſi manu tranſverſus forti-
er conſtringeretur, noctu præcipuè urget, dum
eger lecto incaleſcit. Curvatur penis à fræno
contracto. Accedit ardor urinæ, inter mingen-
dum vix perceptibilis, ſed mictu finito, per
urethræ

urethtæ ductum acriter urens, maximè quà in glandem desinit. Caruncułæ lotii exitum quandoque impediunt; & scroti dolor ac inflammatio nonnunquam superveniunt.

℞ *Mass. Pil. Coch. maj.* ℥iij. *Extract. Rud.* ℥j. *Resin. Jalap. & Diagryd. an.* ℥ß. *cum s.q. Opobalsam. f. Pilul. è singulis drachmis vi,* cap. *iv.* singulis Auroris, donec urinæ ardor & color flavus materiæ seminiformis, admodum imminuantur. Posteà alternis diebus ad septimanas adhuc duas; ac sub finem bis in septimana, donec humor stillare prorsus cessaverit.

In iis, qui difficulter purgantur, subinde intercalari possit potio communis purgativa, cui addantur, *Syr. è Spin. Cervin. & Elect. Succ. Rosar. an.* ℥ij. vel, curatione minùs fœliciter succedente *Turbith. Minerale ad gr. viij.* exhiberi potest bis vel ter, diebus quatuor interpositis, vel vice *Turbith. Mineral.*

℞ *Pil. ex duob.* ℥ß. *Merc. dulc.* ℈j. *cum s. q. Opobalsam. f. Pil. iv.* summo mane sumendæ.

Si æger à dicta Catharsi abhorreat, post pilulas primò descriptas, tribus Auroris continuis sumptas, tum manè & horâ quintâ pomeridianâ quotidie injiciatur Clyster sequens, nisi quod semel aut bis in hebdomade, omisso Clystere, Cathartic. imperandum.

℞ E-

℞ *Elect. è ſucc. Roſar.* ℥vj. *Terebinth. Venet. Vitello ovi ſolut.* ℥ſs. *diſſolvantur in Aq. hord.* ℔j. *Colatur. adde Syrup. Violar.* ℥ij. *M. f. Clyſter.*

Singulis noctibus cap. *Opobalſam. vel Balſam. è Mech. gutt. xxv. Sacch. fruſtulo exceptas.* Hujus defectu *Terebinth. Cypr.* ad quantitatem Avellanæ ſubſtitui poſſit.

Sit pro potu hydrogala; at Cereviſia tenuis in paſtibus.

Vel ſequentem methodum aggrediatur,

℞ *Pil. ex duob.* ℥ſs. *Opobalſam. gutt. iij. f. Pil.* N° *iij.* Cap. horâ quartâ matutinâ, ſuperdormiendo, repetanturque alternis diebus, vel tertiâ quâque die.

Diebus à purgatione vacuis injiciatur manè & horâ quintâ pomeridianâ Clyſter ſupra deſcriptus.

Procedente curatione (quâcunque eâ demum methodo tentetur) ſanguis ſub medium curationis ſemel vel bis mittatur, & diæta refrigerans & incraſſans præſcribatur, & medicamenta iiſdem virtutibus pollentia, qualia ſunt *hydrogala, Emulſiones ex ſeminib. frigid. &c.*

Si Penis Tumore afficiatur,

℞ *Rad. Althææ, Lilior. an.* ℥iſs. *fol. Malv. Verbaſc. Sambuc. Hyoſcyam. flor. Chamamel. Melilot. an. M. j. ſem. Lini, Fænugræc. an.* ℥ſs. *coq. ſ. q. Aq. font. & f. fotus* parti affectæ admovendus per horæ unius ſpatium bis vel ter in die.

Per-

Peracta fomentatione, perungatur pars affecta ol. ſem. Lin. rec. extract. dein *Emplaſt. è mucilaginibus aluta agninæ inductum* labiis præputii tumidis circumponatur.

Si Ulcus in præputii labiis, vel in ſubjecta Glande, hæreat, tum ſequens Linimentum præſcribatur.

℞ *Unguent. Baſilic.* ℥vj. *Unguent. Nicotian.* ℥ij. *præcipitat. Aq. Roſar. loti & optimè levigat,* ℥ſs. *M. f. linimentum,* quo lintei raſura imbuatur, & ulcuſculo admoveatur ſemper poſt fotûs præſcripti uſum.

Si ſcrotum tumore afficiatur, ſtatim ſanguis è brachio extrahatur, atque bis in die fotus præſcriptus parti affectæ admoveatur, ſingul. fotibus addendo *Sp. vin. Gallic. Cochl. j. vel alterum, vel Cataplaſma ex Oxycrato & farina fabarum* ejus loco applicandum.

Catharticis interim & refrigerantibus modò deſcriptis introſumptis utendum eſt.

## *De Lue Venerea.*

Ubi aut à Gonorrhœæ mora, aut indebito aſtringentium uſu labes ſanguini imprimitur, exoritur Lues. Comparent in inguine bubones, in capite dolores & artibus, articulorum interſtitiis, noctu præcipuè invadentes in lecto calidos, cruſtulæ & porrigines in variis corporis partibus, quæ ad inſtar favi flaveſcunt. Quo magis

magis ſcabies diffunditur, tanto minùs æger diſcruciatur. In cranio, tibiis barchiiſque exoſtoſes, oſſium inflammatio & caries; ulcera phagædenica in variis partibus, ſed fauces utplurimum primò aggredientia, quæ ſenſim per palatum in naſi cartilaginem propagata mox eam abſumunt, ita ut fulcro deſtitutus naſus planè concidat. Ulceribus, carie & dolore indies exacerbatis, pereunt per vices membra; lacerum donec cadaver, & jam Superis grave humo condatur.

℞ *Axung. Porc.* ℥ij. *Mercur. crud.* ℥ß. *M. f. Linimentum*, in tres æquales partes dividendum, quarum unâ brachia, crura & tibias ad tres noctes continuas manibus ſuis æger perungat.

Si intra tres dies poſt ultimam inunctionem ſalivatio haud procedat, tum *gr. viij. Turbith. mineral. in Conſerv. Roſar. rubr. exhibeantur; aut corporibus minus robuſtis, Merc. dulc.* ℥ß. Sed ſi ſalivatio jam incepta priùs elanguſcat, quàm ſymptomata penitus diſparuerint, extimulanda ſubindè eſt eâdem doſi Mercur. dulc.

Ptyaliſmus ita moderari debet, ut ℔iv. Nycthemeri ſpatio expuat, aut circiter.

Si juſto altiùs exurgat Ptyaliſmus, cum inflammatione nimia, atque id genus ſymptomatis ingruentibus, aliquantulum reprimendus eſt medicamentis Catharticis, donec debitum gradum attingat.

Quamprimum ſymptomata ceſſaverint, ſtatim immutentur induſium & tori lintea; at ſint ea, quæ priùs uſurpata fuerint.

Si Diarrhœa ſupervenerit (quod plerunque accidit antequam ſalivatio bene aſſurgat) tunc ſiſtenda eſt uſu Laudan. Liquid. ita doſin augendo atque repetendo, ut debitum opus abſolvat.

Si os exulceretur, *Aq. Roſ. Damaſcen. vel lacte Aq. permiſto*, vel ſeq. decoct. idem colluatur.

℞ *Rad. Alth. Hord. mundat. an.* ℥*j. ſem. Cydonior.* ℥*ſs. coq. ſ. q. aq. ad* ℔*ij. M. f. Gargariſma*, quo ſæpius utatur.

Diæta & reliquum ægri regimen, eadem ac in Catharſi imperanda, niſi quòd primis aliquot diebus liquore poſſetico, juſculis avenac. hordeac. ac cereviſiâ tenui tepefactâ nutriatur.

His omnibus ritè peractis (etiamſi ſymptomata omnia evanuerint, ac ſubindè morbus penitus profligari videatur) quo minùs recidivam poſtliminiò patiatur, ægro ſemel in ſeptimana per 4 vel 6 vices, *Merc. dulc.* ℥*j.* exhibendus, utcunque bene ſe habeat, & etiamſi jam domo egrediatur.

## *De Fluore albo.*

Quod effluit, modò album, modò pallidum, flavum, viride aut ſubnigrum, modò acre & ero-

erodens, tetrum quandoque & graveolens, decoloratur facies, dolet ſpina dorſi, proſternitur appetentia, oculi pedeſque intumeſcunt.

Mittatur ſanguis è brachio ad ℥ viij.

℞ *Pil. Coch. maj.* ℈ij. *Caſtor. gr. ij. Balſam. Peruv. gutt. iij. f. Pil.* N° *iv.* cap. horâ quartâ matutinâ, ſuperdormiendo; repetanturque per duas alias vices interponendo diem unum vel alterum pro ratione virium.

℞ *Aq. Rut.* ℥ *iv. Aq. Bryon. comp.* ℥ *ij. Sacchar. Cryſtal. q. ſ. f. Julapium*; Cap. Cochl. *iij.* vel *iv.* in omni languore; deinde aggrediatur uſum ſequentium.

℞ *Theriac. Andromac.* ℥ *iſs. Conſerv. flaved. Aurant.* ℥ *j. Diaſcord.* ℥ *ſs. Zinzib. condit. & nuc. moſchat. condit. an.* ℥ *iij. Pulv. è Chel. cancr. comp.* ℥ *iſs. cort. exterior. granat. Rad. Angelic. Hiſpan. corall. rubr. præparat. & Trochiſc. de terra Lemnia an.* ℥ *j. Boli Armen.* ℈ *ij. Gumm. Arabic.* ℥ *ſs. cum ſ. q. Syr. de Roſis ſiccis f. Electuar.* de quo ſumat ad magnitud. nuc. moſchat. maj. manè, horâ quintâ pomeridianâ, & nocte per menſem integrum, ſuperbibendo ſequent. infuſionis Cochl. *vj.*

℞ *Rad. Enul. campan. Imperator. Angelic. Calami Aromat. an.* ℥ *ſs. fol. Abſinth. Roman. Marrub. alb. Centaur. min. Calamynth. vulgar.*

*Sulv. ſicc. an. M. j. Bac. Junip. ℥ij. incidantur minutim & infundantur in Vini Canarini* ℔iv. *ſtent ſimul in infuſione frigida*, coletur tantummodò tempore uſûs.

Veſcatur cibis eupeptis, & abſtineat ab uſu olerum & fructuum qualiumcunque, & ſingul. paſtibus bibat vinum Hiſpanicum.

*De Diabete.*

Succi ſanguini illati per vias urinarias crudi & inconcocti exitum ſibi quærunt, unde ſenſim labefactantur vires, colliqueſcit corpus, & quaſi ſubſtantia ejus per hanc cloacam exinanitur, cum ſiti, ardore viſcerum, lumborum coxarumque intumeſcentiâ, & ſalivæ ſpumoſæ expuitione crebrâ.

Eadem præſcribenda ſunt, quæ in fluore albo, omiſſis ſolummodò Venæſectione & Catharſi.

*De Dolore Hæmorrhoidum.*

Intenſiſſimus inter excernendum dolor excruciat, & excrementi ſuperficies cruore conſpergitur: Nonnunquam tumores verrucis ſimiles in ſphinctere deliteſcunt, vel in margine ani apparent.

Mittatur ſanguis è brachio dextro ad ℥x.

℞ *Sem. Melon. Pepon. an.* ℥ſſ. *ſem. Papav. alb.* ℥ij. *Amygd. dulc. exemptis.* N°v; *contundantur*

*tundantur in mortario marmoreo, ſenſim affundend. Aq. Hord.* ℔ *iſſ. Aq. Roſar.* ℥ *ij. Sacchar. albiſſ. q. ſ. f. ſ. a.* emulſio; cap. ℥ *iij.* ſæpiuſcule.

℞ *Flor. Sulphuris, Pulv. Glycyrrhiz. Salviæ an.* ℥ *j. cum ſ. q. Balſam. Lucatel. f. Pil. è ſing.* ℈ *vij.* cap. *iij.* ter in die, ſuperbibend. Emulſ. præſcript. Cochl. *vj.*

℞ *Aq. Spermat. Ranar.* ℥ *iv. in qua diſſolv. Lithargyr.* ℥ *ij. Opii* ℈ *j. f. mixtur.* cujus tantillo Linteum madefactum parti affectæ applicetur, vel ſi tumor intus lateat, Cochl. *iij.* ejuſdem mixturæ modo Clyſteris injiciantur.

Pro Potu ordinario bibat aquam cum lacte coctam, vel aquam hordei; & à carnis uſu abſtineat. Singulis noctibus ſumat Diacodium.

### *De fluxu Hæmorrhoidum immodico.*

Innoteſcit ex viribus proſtratis, copia & diuturnitate ſanguinis excreti, fœdo corporis colore, ad citrinum (ut in Ictero) vergente. Succedit cachexia, & deinde hydrops.

Eadem conferunt, quæ in fluxu menſium immodico, omiſsâ ſolummodò Catharſi.

### *De Epilepſia Puerorum.*

Vel primo menſe aggreditur à frequentibus nimis alvi dejectionibus (quo in caſu *Diaſcor-*

*dium ad quantitatem grani piperis in Aq. Saxifrag. aut lacte materno ſolutum*, & exhibitum præſtantiſſimum eſt remedium.) Vel circa dentitionis tempus, à ſeptimo ad decimum menſem, tuſſi comitata eſt, aut (quod multo deterius) vomitu & diarrhœâ, viridia (ut ſolent hyſtericæ) egerente infante. Quandoque improviſò occupat paroxyſmus, os oculóſque diſtorquens, nigredine vultum perfundens, & artus varios covellens; quandoque præſentitur ex digitorum in pugnum contractione, & ex ſtabili & inſolita oculorum orbitarum immobilitate. Nunc ſeriùs, nunc ocyùs ſe propipiunt paroxyſmi, certâ ſubinde periodo, vago aliàs inſultu revertentes; ſed, appetente morte, à tergo ſe invicem premunt: Si quando inducias dent, in ſoporem proclives ſunt infantes, donec à novo reſuſcitentur inſultu.

Applicetur *Emplaſtr. Epiſpaſtic.* cervici quamprimum.

℞ *Aq. Epilept. Lang.* ℥iij. *Laudan. Liquid. gutt. j. vel ij. vel plures, pro ratione ætatis, Syr. Pæon.* ℥j. *M. f. Hauſtus*, ſumendus quamprimum.

℞ *Aq. Rutæ* ℥iij. *Aq. Epilep. Lang. & Aq. Bryon. comp. an.* ℥j. *Syr. Caryophyll.* ℥ſs. *M. f. Julap.* cap. Cochl. *j.* ſingulis horis, ſi hauſtus præſcript. paroxyſmum non fugaverit.

*De Rachitide.*

In Rachitide adſunt partium laxitas & mollities, debilitas & languor, ignavia & torpor, & inæqualis earundem nutritio, *viz.* caput juſto majus, facies plenior & floridior, partes carnoſæ infra caput emaciantur, nodis protuberant articuli, maximè carpi, paulo minùs tali, tument & extremitates coſtarum, oſſa incurvantur, præcipuè tibiæ & fibulæ, hinc ulnæ & radii, aliquando femoris & humeri, dentes tardè & moleſtè erumpunt, vacillant, nigreſcunt & fruſtulatim excidunt, pectus à lateribus anguſtum, anteriùs acuminatum, abdominis repletio, & hypochondriorum tenſio, tuſſis & pulmonum vitia laceſſunt, lateralis decubitûs modò dextri, modò ſiniſtri averſatio.

℞ *Fol. Abſinth. vulgar. Centaur. minor. Marrub. alb. Chamædryos, Scordii, Calamenth. vulg. Parthen. Saxifrag. pratenſ. Hyper. Virg. aureæ, Serpill. Menth. Salv. Rutæ, Card. Benedict. Puleg. Abrotan. Chamæmel. Tanacet. Lilior. convall. omnium rec. collect. & inciſ. an. M. j. axung. Porcin.* ℔iv. *ſev. ovin. & Vin. claret. an.* ℔ij. *macerentur in olla fictili ſuper cineres calides per horas xij. deinde ebulliant ad humiditatis conſumptionem, & poſtea colentur, ut f. Linimentum,* quo venter & hypochondria illinantur

manè & ſerò, ut etiam & artus hoc malo affecti, per 30 vel 40 dies, vel donec convaluerit.

℞ *Singulor. ingredient. præſcript. an. M. ij.* infundantur frigidè in Cereviſiæ non Lupulat. Cong. *iv.* pro potu.

In Tumoribus Infantum ventres occupantibus ab Evacuationibus nimiis procedentibus,

Sanguis & Viſcera herbis corroborantibus confortentur, ut in Rachitide vera; niſi quod axillæ etiam utræque inungantur, at non artus.

## *De Febribus à Dentitione Ortis.*

℞ *C. C. gutt. ij. vel iij. vel iv. pro ratione* ætatis in Cochl. *j.* vel altero Aq. Ceraſor. nigr. vel Julap. alterius appropriati, exhibeantur quartâ quâque horâ ad vices quatuor, vel ſex.

## *De Febre hectica Infantum.*

Sine inſigni calore languent cum appetitûs dejectione, artus & truncus corporis emaciantur.

℞ *Rhabarb. inciſi* ℥*ij. in lagenam vitream cereviſiæ tenuis* ℔*ij. capacem* immittantur pro potu ordinario

Epoto

Epoto hoc liquore, *alia* ℔ij. *eidem Rhabarb. ſuperaffundantur ad tertiam vicem.*

Si verò potus vi Catharticâ ſaturatius imprægnetur, tum ℔j. epotâ alteri cereviſia mox adjiciatur.

### *De Tuſſi Puerorum Convulſiva.*

Phlebotomiâ, & repetitâ Catharſi, eâque ſolâ, expugnatur hæc tuſſis, malum aliàs pertinaciſſimum, & fere inſuperabile; ſed leniora tantum cathartica hic conveniunt, eaque cochleatim ingerenda ſunt, pro ratione ætatis.

### *De Hæmorrhagia Narium.*

Dolor & calor ſinciput lancinant.

Sanguis è brachio frequenter extrahatur; victûs ratio refrigerans & incraſſans injungatur; Julap. refriger. & incraſſans, cum Emulſionib. refrigerantib. præſcribantur.

Singulis diebus injiciatur *Enema refrigerans*, atque omni nocte *Paregoric. ex Syrup. de Mecon.* exhibeatur, & ſemel vel bis *potio purgativa commun.* præſcribatur.

*Linteа quadruplicata immerſa Aq. frigidæ, in quâ ſal Prunell. diſſolvatur, & poſtea leviter expreſſa* admoveantur nuchæ, & utrinque cervici ſæpius in die.

Poſt Evacuationes liquor ſeq. applicetur.

℞ *Vitriol. Hungaric. & Alum. an.* ℥j. *Phlegmat. Vitriol.* ℔ſs. *Coq. tamdiu, donec omnia fuerint diſſoluta, liquorem frigefactum filtra, & à Cryſtallis ſubinde natis ſepara, liquori reſiduo adde Ol. Vitrioli duodecimam partem*; vel potiùs

℞ *Aq. Plantag.* ℥iij. *Bol. Armen. ſubtiliſſ. pulver.* ℥ſs. *probè miſceantur*; *Turunda ex linteo raſo liquore hoc humectata*, nari, ex quo ſanguis ſtillat, indatur per dies duos iſtic relinquenda.

Hoc minus ſuccedente, *Vitriol. Rom. Aq. commun. diſſolvatur, qua Turunda imbuta* nari indatur, *liquore* etiam *hoc linteamina imbuta*, & applicata ſanguinem ſiſtunt ſi ex parte externa diffluat.

## *De Chloroſi.*

Adſunt vultûs & totius corporis decoloratio, intumeſcentia in facie, palpebris & malleolis, gravitas totius corporis, crurum ac pedum tenſi a laſſitudo, reſpiratio difficilis, cordis palpitatio, capitis dolor, pulſus febrilis, ſomnolentia, pica & menſium ſuppreſſio.

Sumat

Sumat *Pilulas Chalybeatas*, vel *Pulv. Chalybeatum* in affectione hyſterica præſcript. plus minus pro ratione ætatis, ſuperbibend. hauſtum vini cujuſlibet ad libitum, vel infuſion. corroborant. cum Rad. Angelic. ibidem deſcriptam. Si Ægra non valde debilis fuerit, purgetur ſemel vel bis ante præſcriptorum uſum.

### *De Suppreſſione Menſium.*

Cibum faſtidiunt mulierculæ, fœdus faciei color eſt, gravitas totius corporis, dolores ſincipitis, lumborum, crurum & imi ventris; pedum tumores infeſtant.

Eadem præſcribenda ac in Hyſterica affectione; at ſi iiſdem remediis affectus non ceſſerit, tum ſequ. ſunt præſcribenda.

Singulis Auroris, & quartâ pomeridianâ, ſumat *Julap. Hyſteric. ſine Caſtor. Cochl. v. inſtilland. Sp. C. C. gutt. xij.* Singulis noctibus ſumat cubitum itura *Trochiſc. de Myrr.* ℈ß. *cum Syr. de Artemiſ. in formam boli vel Pil. redact.*

### *De Vomitu ſanguinis, vel etiam de Hæmoptoë.*

In Hæmoptoë dolor & calor pectus obſident, cum debilitate quadam.

Mittatur

Mittatur ſanguis ex brachio dextro ad ℥x. Sequenti manè ſumat *potionem purgativ. communem.* & eadem nocte *Paregoric. ex* ℥iij. *Aq. Cinaſar. nigr. & Syr. de Mecon.* ℥j.

Poſtea aggrediatur uſum ſequent.

℞ *Bol. Armen.* ℥j. *Pulv. Radic. Symphyt.* ℥ij. *Ter. ſigillat. Lapid. hæmatit. & Sang. Dracon. an.* ℥j. *Sacchar. albiſſ. ad pondus omnium. M. f. Pulvis ſubtiliſſimus,* cujus cap. ℥j. manè, quintâ pomeridianâ, & nocte, ſuperbibend. ſequent. *Apozemat.* ℥iv.

℞ *Fol. Plantag. Rubi Sylveſtris, & Millefol. an. M. j. coq. ſ. q. Aq. ad* ℔iſs. *in Colatura diſſolv. Syrup. de Symphyt.* ℥ij: *M. f. Apozema*; Vel cap. Cochl. *vj.* ſequent. Tincturæ.

℞ *Flor. Roſ. rubr.* ℥vj. *Cort. intern. querc.* ℥ſs. *ſem. Plantag. parùm contuſ.* ℥iij. *Aq. font.* ℔ij. *Sp. Vitriol. q. ſ. ad levem aciditatem, infundantur in vaſe clauſo, calore leni, per 4 horas*; *Colatur. adde Aq. cinnam. hordeat.* ℥iij. *Sacchar. albiſſ. q. ſ. ut f. Tinctura guſtui grata.*

Si ab uſu Pulveris abhorreat, utatur Electuario in fluxu menſium immodico præſcripto.

Enema quotidie injiciatur, & Diacodium horâ ſomni propinetur.

Phle-

Phlebotomia inſuper ſemel, bis, vel ter (ut opus fuerit) reiteretur, diebus aliquot interpoſitis; & ſemel iterum, ſi res poſtulet, repetatur potio purgativa.

Diæta incraſſans & refrigerans obſervanda.

*De Punctura Tendinis.*

Humor aqueus, vel ichor quidam, ex orificio venæ inciſæ jugiter deſtillat.

℞ *Rad. Lilior. alb. in lacte vaccin. ad teneritud. coct. & contuſ.* ℥iv. *farin. ſem. Lin. & Avenac. aa.* ℥iij. *coq. ad cataplaſmatis conſiſtentiam in eadem lacte, quo radices coctæ fuere*, applicetur parti affectæ manè & ſeró.

*De Ambuſtis.*

Foveatur pars affecta *linteis Sp. Vini imbutis*, uſque dum evaneſcat dolor, ac deinceps de novo imbuatur *linteum*, quod ter vel quater in die parti affectæ applicetur.

*De Mania communi.*

A nimis exaltata & vivida ſanguinis craſi oritur. Eſt & alia maniæ ſpecies febres intermittentes diuturniores excipiens, in ſtultitiam tandem

tandem degenerans, producta quidem à debilitate & vappiditate ſanguinis, à diutina fermentatione. Præſcribantur itaque cardiaca generoſiora, *viz. Theriac. Andromach. Electuar. de ovo, Pulv. Comitiſſ. Pulv. Gualt. Rawleigh, in Aq. Epidem. Theriacal. vel aliâ*; & diæta inſtituatur analeptica.

In junioribus mittatur ſanguis è brachio ad ℥ *viij.* vel *ix.* bis vel ter, diebus tribus inter ſingulas Venæſectiones interpoſitis. Tum ſemel extrahatur ſanguis è venis jugularibus. Poſtea tota curatio innitatur ſequenti Cathartico, quod ſumat tertiâ vel quartâ quaque die, donec convaluerit; hoc interim obſervando, quod cùm octies vel decies purgatus fuerit æger, exhibitio Cathartici ad ſeptimanam unam vel alteram omittatur.

℞ *Pharmac. domeſtic.* (i. e. *Rad. Bryon. alb.*) *pulv.* ℥ *j.* lactis vaccin. ℥ *iv.* vel,

℞ *Ejuſdem* ℥ *ſs. vel* ℥ *vj. Vin. alb.* ℥ *iv. infundantur per noctem, in colatura diſſolv. Syr. Violar.* ℥ *j. M. f. Potio*; vel,

℞ *Gombogii gr. xiv. Aq. Ceraſor. nigr.* ℥ *iij. Syr. Caryophyll.* ℥ *ſs. f. Potio.*

De

*De Contuſionibus.*

Mittatur ſanguis è brachio lateris magis affecti ad ℥x. ſeq. manè exhibeatur *potio communis purgativ.* ac deinceps eodem modo Venæſectio & purgatio alternatim repetantur, donec æger convaluerit. Interim toto decurſu, ſi internæ partes lædantur, ſequentia præſcribantur.

℞ *Decoct. Pectoral.* ℔iſs. *Syr. Violar. cap. Ven. an.* ℥ij. *f. Apozema*; cap. ℔ſs. ter in die. Item *Ol. Amygd. dulc. rec. Cochl. j.* ſæpius propinetur.

℞ *Ol. Amygd. dulc. unguent. Dialth. & Pomat. an.* ℥j. *M. f. Linimentum*, quo ungatur pars læſa manè & ſerò, ſuperimponed. fol. Braſſicæ.

*De furfure ſive porrigine capitis.*

*Potio purgans communis* bis propinetur, dein

℞ *Ol. Amygd. Amarar. & Laurini, Cinorum fol. Abrotani an.* ℥j. *miſceantur probè, & f. Linimentum*, quo inungatur totum caput ſingulis matutinis, diligenter infricando, & poſtea ſuperimponendo veſicam ſuillam.

At

At primùm abradatur totum capillitium, & poſtea defricentur ſquammulæ ſenſim ſingulis auroris, prout decidere pronæ fuerint.

*De Morſu canis Rabidi.*

Poſt 40 dies aut plures melancholica adſunt ſymptomata, febris, ſitis, hydrophobia, & tandem partium extremarum convulſio.

℞ *Sp. vin. optimè rectificat.* ℥iv. *Theriac. Andromach.* ℥j. *f. mixtura,* qua illinatur pars læſa ter in die, ſuperimponendo linteum eâdem mixturâ imbutum.

*De Ulcere Veſicæ.*

Pus graveolens, aut ſanguis nonnunquam ſquammulæ ſeu pelliculæ membranaceæ ac veluti furfuraceæ cruſtulæ (ſin in renibus hæreat ulcus, carunculæ minutulæ, aliquando craſſiores) unà cum urina excernuntur, perpetua dyſuria & continuus earum partium dolor infeſtant; renibus autem affectis ulcere, dilucida dyſuriæ, & doloris dantur intervalla, pus etiam excretum copioſius eſt, album, leve & minimè fœtidum, lacti ſimilis profunditur urina: & longè poſt mictionem pus ab eo ſeparatum in fundo reſidet.

℞ *Em-*

℞ *Emplaſt. Flor. unguent. dict.* ℥iſs. *f. Pil.* N° *ix.* cap. *iij.* horis medicinalibus ſuperobibendo ſequentis Aq. Cochl. *vj.*

℞ *Rad. Fœnicul. Conſol. Ariſtoloch. & Caryophyll. an.* ℥iij. *fol. Agrimon. Hyper. Bugul. Sanicul. Plantag. an. m. vj. incidantur minutim, & diſtillentur in vini alb. & lact. an.* ℔iv. *& eliciantur ſolummodo* ℔iv. pro uſu.

℞ *Pulv. Rad. Conſolid. maj. & Gumm. Arabic. an.* ℥j. *Sacchar. Penid.* ℥iſs. *f. Pulv.* cap. bis in die.

*De Aſthmate inveterato in iis, qui ſanguineo temperamento praditi ſunt.*

Dividitur in tres ſpecies; Dyſpnœam, quæ difficilis eſt reſpiratio, in qua crebrior denſiorque eſt ſpiritûs tractio ex infarctu pulmonum ſine ſtertore oriunda: Aſthma ubi frequens & magna reſpiratio in qua diaphragma, muſculi intercoſtales, tum etiam abdominis, violenter moventur, cum ſtertore & ſibilo. In priore ſpecie ſubſtantia pulmonum, in poſteriore bronchia oppilantur: Et Orthopnœam, quæ eſt maxima ſpirandi difficultas, ſub qua ægrotantes non niſi ſedentes, & erecta cervice reſpirare poſſunt. Muſculi etiam pectoris & ſcapularum vehementer exagitantur.

Mitta-

Mittatur ſanguis è brachio dextro ad ℥x. Die ſequenti ſumat *potionem purgativam communem*, quæ repetatur tertiâ quâque die per duas alias vices.

Diebus à Purgatione vacuis ſumat ſequentia.

℞ *Sem. Aniſ. ſubtiliſſ. pulv.* ℈ij. *cum ſ. q. Balſam. Lucatell. f. pil. ex ſingulis drachmis vj.* cap. *iij.* manè & quintâ pomeridianâ, ſuperbibend. *decocti Amari ſine purgantibus* ℥iv. calidè.

Si Symptomata adhuc maneant, repetatur de novo integer Proceſſus.

## *De Paralyſi.*

Senſus & motus in partibus quæ afficiuntur, vel ſimul, vel alterutra tantùm facultas aut prorſus extinguitur aut vitiatur.

Sumat *pil. Coch. min.* ℈ij. alternis matutinis per *vj.* vices; deinde capiat ter in die per *xxx.* dies *Electuar. Antiſcorbutic.* ℥ij. ſuperbibend. *Aq. Antiſcorbutic. Cochl. vj.* Vide in Scorbuto.

℞ *Unguent. Nervin.* ℥iij. *Sp. Lavendul. comp. & Sp. Cochl. hortenſ. an.* ℥iſs. *M.* & inungantur partes affectæ, ut ſpina dorſi, &c. manè & ſerò.

Plera-

Pleraque præſcriptorum, utut curationi ſcorbuti ſolùm deſtinari videantur; attamen cùm humores crudos ac fixos volatiliſare præ aliis apta nata ſunt, in Paralyſi etiam conveniunt.

## *De Tuſſi & Phthiſi.*

Tuſſis per ſe patet. Phthiſis autem invadit ab ætatis anno 18 ad 35. Extenuatur totum corpus. Febris moleſtat hectica, à cibo invaleſcens (quæ ex celeritate pulsûs, rubedinéque genarum dignoſcitur) materia per tuſſim excreta cruenta eſt aut purulenta; in ignem conjecta tetrum ſpirat odorem, & immiſſa in vas aquâ repletum, fundum petit. Sudores nocturni ſuperveniunt, liveſcunt tandem genæ, facies pallet, exacuantur nares; temporum quoque collapſus, unguium incurvatio, capillorum defluxio, & alvi profluvium colliquativum, mortem ſtatim ſubſecuturam denunciant.

Si Tuſſis recens fuerit, & nondum febrim, aut alia ſigna Peripneumoniam notham comitantia, accerſiverit, vel ſi à Pleuritide, aut Peripneumoniâ, ob neglectam ſufficientem Phlebotomiam, malè curatis eadem nequaquam orta fuerit, ſatis erit, ut æger per dies aliquot ab uſu vini & carnis abſtineat, & his ſimilibus remediis ad libitum utatur.

*Balſa-*

*Balſamum ſulphuris miſtum ad gutt. x. Cochl. j. Sacchar. cryſtall. inſtillatum* bis vel ter in die devoretur; ut etiam Tabellar. ſequent. quas ſecum æger continuò portet, unam frequenter aſſumat.

℞ *Sacchar. cand.* ℔ij. *coq. ſ. q. Aq. font. uſque dum adhæreſcat extremis digitor. tum adde pulv. Glycyrrhizæ, Enul. Campan. ſem. Aniſ. & Angelic. an.* ℥ſs. *pulv. Irid. Florentin. Sulphur. an.* ℥ij. *Ol. Chym. Aniſi* ℥ij. *f. ſ. a. Tabellæ domeſticæ vocand.* atque hoc interim Eclegmate utatur.

℞ *Ol. Amygd. dulc.* ℥iij. *Syr. cap. Ven. Violac. an.* ℥j. *Sacchar. cand. q. ſ. f. Lohoch.* de quo ſæpius lambat bacillo Glycyrrhizæ, quando tuſſis urget.

In defluxione tenui *Eclegmata inoraſſantia* exhiberi poterint.

Sin verò Tuſſis his remediis haud ceſſaret, (præcipuè autem ſi febrim comitem habeat, vel ſi à Peripneumonia aut Pleuritide originem ducat) tum ineptum fuerit pectoralibus fidere; ſed Veneſectione & Catharſi debellanda eſt, ut in methodo, quæ in curatione Peripneumoniæ nothæ deſcribitur.

Verùmenimverò cùm, his omnibus non obſtantibus, Tuſſis non modò non curationi ceſſerit, ſed & diuturnâ Pulmonum concuſſione eos

eos adeò debilitaverit, ut Phthisis tandem subingressa fuerit, tum hâc methodo procedendum.

℞ *Balsam. Peruv. gutt. x. instillentur Cochl. j. Syr. Hederæ terrestris*, vel si id minus gratum fuerit ægro, *Coch. j. Sacchar. crystal.* cap. ter in die, superbibendo *Decocti Amari sine purgantib.* ℥iv. vel (si idem ventrem moverit) ℥iij.

Sed omnibus aliis (quotquot adhuc inventa sunt) *Equitatio* ad satis longa & diutina itinera facilè palmam præripit, hoc observando, ut si æger juvenilem ætatem prætergressus fuerit, plus temporis huic exercitio impendere debet, quàm si puer aut juvenis esset, & sanè haud multò certiùs Cortex Peruv. febri intermittenti, quàm in hac ætate Equitatio Phthisi medetur.

## *De Scorbuto.*

Adsunt spontaneæ lassitudines, corporis gravitas, dyspnœa, præcipuè ex motu corporis, gingivarum putrefactio, oris graveolentia, sanguinis è naribus crebra eruptio, ambulandi difficultas, crurum nunc intumescentia, nunc extenuatio, in iisdem perpetuo maculæ lividæ, plumbeæ, flavæ, aut violaceæ, faciei color plerumque ex pallido fuscus.

Ex-

Extrahatur ſanguis è brachio ad ℥ viij. niſi ſigna Hydropis adfuerint: Seq. mane exhibeatur Potio Purgat. communis, quæ repetatur tertiâ quâque die per duas alias vices.

Diebus à purgatione vacuis, ut etiam poſtea, per menſem unum vel alterum ſumat ſequentia.

℞ *Conſerv. Cochl. hortenſ.* ℥ ij. *Conſerv. Lujul.* ℥ j. *pulv. Ari compoſ.* ℥ vj. *cum ſ. q. Syr. Aurantior. f. Electuar.* de quo cap. ad magnitud. nuc. moſchat. maj. manè, quintâ pomeridianâ & nocte, ſuperbibendo *Aq. Raphan. compoſ. Cochl.* vj. vel ſequentis.

℞ *Rad. Raphan. Ruſtican. raſpat.* ℔ ij. *rad. Ari* ℔ j. *fol. Cochl. hort. M. xij. fol. Menth. Salv. Naſturt. aquat. & Becabung. an. M.* vj. *Sem. Cochl. hort. parùm contuſ.* ℔ ſs. *nuc. Moſchat.* ℥ ſs. *Vin. alb.* ℔ xij. *diſtillentur organis communib. & eliciantur ſolummodo* ℔ vj. pro uſu.

Vel *Folia rec. Cochl. hortenſ. per ſe diſtillentur* pro uſu dicto.

Porrò ſequens cereviſia ſit pro potu ordinario.

℞ *Rad. Raphan. Ruſtican. recentis inciſi* ℥ ij. *fol. Cochl. hort.* N° xij. *paſſular. enucleat.* N° vj. *Aurant. cum toto inciſi dimidium, in lagenam vitream* ℔ ij. *Cereviſia tenuis capacem ſubere mox probè obturandam immittantur.*

In

In hunc modum parentur eodem tempore ſex Lagenæ pro uſu, & poſt aliquot dies ſex aliæ, antequam priores exhauriantur, & ſic deinceps; vel,

Vice hujus Cereviſiæ ſeq. mixturæ Cochl. *iij.* vel *iv.* immiſceri poterint ſingulis hauſtibus potûs communis.

℞ *Rad. Raphan. Ruſtican. & ſem. Cochl. hortenſ. an.* ℥*ſs. fol. Cochl. hortenſ. M. ij. pulpam unius Aurant. Contundantur ſimul in mortar. marm. ſenſim affundendo vin. alb.* ℔*ſs. Colatur cum levi expreſſione,* & reponatur pro uſu.

Eadem remedia, Phlebotomiâ & Catharſi omiſſis, in Rheumatiſmo ſcorbutico, ſive Hyſterico, valde proſunt.

## *Tabis Deſcriptio & Cura.*

Plures ſunt Tabis ſpecies: Prima ferè oritur è frigore hyeme contracto. Paulo enim ante ſolſtitium Brumale quamplurimi ab ingruente tempeſtate frigidâ, tuſſi tentantur. Quæ cum illos invaſerit quibus pulmones ab ortu ſunt infirmiores, omnino neceſſe eſt ut à repetitis Tuſſiendi conatibus magis adhuc debilitentur; atque ita tandem alimento ſibi proprio aſſimilando, præ mala διαθέσει, prorſus impares reddantur. Hinc jam ingens Pitu-

Pituitæ crudæ copia aggeritur, quæ continuâ pulmonum agitatione & vehementi niſu affatim per eos rejicitur. Pulmones exinde pure referti purulenta μιάσματα quaquaverſum in carnes diſpergunt. Hinc enaſcitur Febris putrida, cujus Paroxyſmus ſub veſperam fatigat, ſub auroram vero ſudore copioſo ac vires imminuente ſolvitur. Denique ad malorum cumulum Diarrhœa ſuccedit, partim ab humoribus putridis ex arteriis Meſentericis in inteſtina depoſitis excretiſque, partim à labefactato peſſundatoque viſcerum Tono. Atque ita demum æger proximâ æſtate fato ſuo fungitur, à morbo, cui prægreſſa Hyems viam ſtraverat, extinctus. Atque hæc eſt præcipua Tabis ſpecies.

Porro cùm ſanguis hac tempeſtate particulis humidis admodum ſcateat, perſpiratioque à ſubitâ pororum contractione plus ſatis inhibeatur, particulæ hæ in pulmones depluunt per interſperſos Venæ Arterioſæ ramos; vel per Ductus Salivales in glandulis fauces occupantibus reponuntur; unde jam humor per Aſperam Arteriam delapſus in Pulmones, Catarrhi ritu, eos ita jugiter irritat, ut creber & vehemens tuſſiendi conatus, tum illam debilitatem, tum alia quæ diximus ſymptomata, brevi accerſat. Cumque non jam amplius naturalem ſibi pulmones œconomiam ſtatumque tueri valeant, Glandulæ & Tubercula adna-

ſci

ſci ſolent; quæ in ægris ex hoc morbo decedentibus, ſi aſpiciantur pulmones, purulentâ ſanie utplurimum diſtenta reperiuntur.

Ubi ſemel invaluerit hic morbus, & radices egerit, remedia plerumque aſpernatur. Tentari utcunque poteſt Medicatio, minuendo cauſam defluxionis in pulmones, pertusâ in Cubito Venâ, Catharticis mitioribus; tum medicamentis pectoralibus dictis, ad varias morbi ſtatus acommodatis, incraſſantibus ſc. ubi humor ſubtilior & tenuior vim Expultricis facultatis videtur eludere; attenuantibus vero ubi pituita craſſior ægre & cum moleſtiâ per os rejicitur. Dein & hectica Febris remediis refrigerantibus contemperantibuſque coercenda. Talia ſunt *Lac Aſininum, Aqua Lactis Stillatitia, Emulſiones ex Amygd. dulc. Sem. Melon. Pepon. Papav. alb. Aq. El. Paralyſ.* &c. Denique & curando Ulceri opem danda eſt; cui rei Terebinthina illa liquida, Opobalſamum vulgo dicta, apprimè conferre conſetur.

Hujuſce Morbi Curatio, me Judice, hâc methodo optimè ac feliciſſimè inſtituetur. Primum educatur è brachio ſanguis, dein ſubducatur Alvus ad tres dies continuos, vel *Pil. Cochli majoribus*, vel *Decocto leniente à nobis præſcripto*. Tertiâ nocte exhibeatur *Syr. de Mec.* ℥ſſ. Intermiſſo biduo, triduove, pro re natâ repetatur Purgatio, idque toties fiat quoties opus eſſe videatur, donec ſymptomata vel

prorſus evanuerint, vel ſaltem miteſcant. Poſt ſingulas Catharſes jam finitas capiat Opobalſami gutt. *xx*. Sacchari particulæ majori inſtillatas permixtaſque, nihil ſuperbibendo. Vel Pil. è Terebinth. ex Inſulâ Chio cum Saccharo cando confectam. Opobalſamum non aſſumendum eſt niſi prægreſſis evacuationibus debitis. Hujus loco *Electuarium è Balſamo Lucatelli, Glycyrrh. pulvere, ſemine Aniſi & Terebinthina* concinnatum poterit ſubſtitui. Poſt evacuationes ſedulo conandum eſt, ut Tuſſis ſedetur, ne Pulmones jugi agitatione infirmiores evadant. Huc facit præ cæteris Syrupus jam laudatus, qui hac formula poſſit uſurpari.

℞ *Decoct. Pectoral.* ℔j. *Syr. de Mecon. Syrup. Capill. Ven. an.* ℥ij. *Cap. Cochl. v. ter in die.* Medicamentum hoc ita frequenter hauſtum catarrhum in Pulmones depluentem paulatim ſiſtet; eoſque in priſtinam ἐυεξίαν reducet, niſi jampridem à naturali ſtatu nimium receſſerint. Promovebit inſuper purulentæ materiæ in iiſdem aggeſtæ concoctionem.

At vero Palmarium hujuſce Morbi præſidium eſt, ut Æger quotidie Equo vehatur. Et qui ſe huic corporis exercitio ſanandum tradit, nullis jam Diætæ legibus adſtrictus, nullo Cibi potuſve genere privandus eſt: cum in hoc uno omnis rei cardo vertatur. Quibuſdam eorum qui ex hac methodo convaluere, tumor

tumor in collo exortus eſt non multum à Scrophulis abludens.

Eſt & alia Tabis ſpecies à Tuſſi prognata, longè diverſo anni tempore, initio ſc. æſtatis. Hæc Juvenes perſæpe illos invadit qui debiliori ſunt corporis habitu, & quorum ſanguis particulis calidis acribuſque refertus eſt. Sanguinem iſti è pulmonibus per tuſſim exantiant, nimio meri potu priùs excalefacti. Ægri dolorem ſive ægritudinem quandam in pulmonibus perſentiſcunt: Quæ Symptomata niſi maturè remediis appropriatis eliminentur, primo ſcabrities, mox & ulcus accedit, unde pus confectum demum expuitur.

Dicta jam Tabis ſpecies Venæſectione & Catharſi bis alternatim repetitis, ſaltem in principio, facile vincetur: Si Diætâ inſuper Refrigerante & Incraſſante utatur æger, & à carnibus in ſolidum abſtineat.

Tertia hujus Morbi Species ortum debet materiæ Febrili in pulmones, Febre jam fatiſcente, amandatæ; qui ſenſim infirmiores redditi à ſymptomatis jam memoratis vexantur.

Quandoque oritur à ſanie iſtâ purulentâ quæ à Pleuriticis non ſatis copiosè per expectorationem egeritur. Quo in caſu eandem quam Empyema, Curationem poſtulat.

Sunt etiam qui à nimis copioſis, & debito frequentioribus evacuationibus debilitati in

quartam Tabis ſpeciem præcipitantur. Hos calor Hecticus à Cibo potuque ſumpto, ſub noctem laceſſit: Atque hos præ cæteris Aphthæ infeſtant.

## *De Podagra.*

Exeunte Januario aut incipiente Februario, ingruit horâ ferè ſecundâ matutinâ, dolore pollicem pedis utplurimum occupante, quandoque vero calcaneum, ſuram aut talum, mox ſequitur rigor cum horrore, & febricitatione aliquali, dolor gradatim intenditur (& pari paſſu rigor atque horror recedunt) idque in ſingulas horas, donec tandem ſub noctem ad apicem pervenerit, nunc tenſionem violentam, nunc morſum canis rodentis, quandoque preſſuram, & coarctationem exprimens; hinc præ exquiſito dolore, nec linteorum ſuperincumbentium pondus, nec cubiculi à fortiori ambulatione concuſſionem ferre Podagricus valeat; dolor non ſedatur, niſi ante horam ſecundam, tertiamve matutinam (decurſo ſcilicet nycthemeri ſpatio à primo appulſu) & jam leni madore perfuſus æger ſomno concedit, à quo evigilans, dolore multum imminuto, partem affectam tumore recenter occupatam cernit. Sequenti die, fortè etiam ad biduum, triduumve dolet nonnihil pars affecta, idque vehementius adveſperaſcente die, levatur autem

ſub

ſub galli cantu. Intra paucos dies alter Pes eodem dolore torquetur. Poſtquam utrumque pedem exercuerit dolor, qui ſequuntur Paroxyſmi abnormes ſunt, tum quoad tempus invaſionis, tum durationem; atque ex ſerie horum paroxymulorum, conſtat Paroxyſmus, qui dicitur Podagræ, diuturnior breviorve pro ægri ætate in iis quæ vegetiores ſunt & quos Podagra rarius invaſit, diebus ſæpe quatuordecim, in ſenioribus atque iis quos ſæpius afflixit, duobus menſibus ferè deſinit, iis verò qui vel annis, vel diuturniore morbi mora fracti ſunt, non prius valedicit, quam æſtas jam adultior eam fugaverit.

Ad quatuordecim primos dies urina coloratior eſt, cum ſedimento rubro, & quaſi arenulis referto, alvus ferè conſtipata eſt. *Appetitus* proſtratio, rigor totius corporis ſub veſperam, integrum Paroxyſmum comitantur, recedentem Paroxyſmum excipit pedis affecti inter digitos pruritus vix ferendus, unde decidunt ſurſures, & pedes deſquamantur.

Hactenus dolor pedibus tantum fuerit infeſtus, cum verò vel ab indebita medicatione turbatur Podagra, vel ob diuturnam morbi pertinaciam, corporis ſubſtantia quaſi in morbi fomitem pervertitur, jam manus, carpos, cubitos, genua, aliaſque regiones occupat; quandoque unum, pluresve digitos diſtorquens, faſciculo radicum paſtinacæ ſimiles eos reddit,

motu paulatim privans, tandemque concretiones, circa articulorum ligamenta, tophaceas generans, cretam, vel oculos cancrorum æmulantes; nonnunquam in cubitis tumorem ſubalbum ciet, ferè ad ovi magnitudinem, qui ſenſim inflammatur cum rubore, nonnunquam femur occupans, ſenſum exhibet quaſi ingentis ponderis appenſi, ſine dolore tamen notabili, at genu exinde petens, id acrius urget, artus jam habet æger ferè undequaque contractos & impeditos, ita ut claudicet, & permoleſte prorepit. Urina illam colore imitatur, quæ in diabete excernitur, dorſum aliæque partes moleſtè pruriunt, maximè ſub horam ſomni.

Poſt multos, diroſque cruciatus Paroxyſmi ſubſequentes (naturâ partim onere materiæ peccantis, partim ſenio oppreſsâ) ægrum quoad dolorem clementius tractant, at loco doloris conſueti, ægritudo quædam cum dolore ventris laſſitudo ſpontanea, quandoque etiam propenſio ad diarrheam ſuperveniunt; quæ ſymptomata evaneſcunt, quoties articulos dolor fatigat, atque ita nunc dolore, nunc ægritudine, quaſi alternatim laceſſentibus, Paroxyſmus longas moras nectit.

Calculum Renum in quamplurimis parit hic affectus.

Fœminas perrarò infeſtat, eaſque, non niſi jam vetulas, & viragines, pueros quoque, & juvenibus minores vix adoritur.

Indi-

Indicatio Curativa in eo vertitur, ut coctioni conſulatur, indigeſtionibus ſublatis, ſeu fit per Pharmaca, ſeu victûs rationem, ſeu exercitationem, ſeu alia ex iſtis quælibet, quæ ſex res non-naturales indigitantur.

Venæſectio, Catharſis, ac Diaphoreſis ſcopum non attingent, licèt iis qui *Renum* ſimul *Calculo*, & exindè orto *Mictu ſanguineo* laborant, non abs re erit, ſemel in ſeptimana, alvum *Mamâ* ſubducere, & ſingulis noctibus, poſt Purgationem Paregoricum exhibere.

Medicamenta, quæ prædictæ indicationi ſatisfaciunt, ea ſunt, quæ moderatè calent, & vel amaricant, vel linguam leniter feriunt, talia ſunt, *Rad. Angelic. Enula, fol. Abſinth. Cent. min. Cham. Chamæp. &c.* quibus addi poſſunt *Antiſcorbutica* dicta, ut, *Rad. Raph. ruſtic. fol. Cochl. hort. Naſturt. aq. &c.* quæ tamen parcius uſurpanda veniunt, quia morbi fomiti aculeos ſubdunt, & calorem augent, præ iſtis aliis, quæ miti calore, atque amaritudine, ventriculum corroborant. Haud inutile eſt *Electuar.* ſequens.

℞ *Conſerv. Cochl. hort.* ℥*iſs. Abſinth. Rom. ſiccor. Aurant. an.* ℥*j. Angel. condit. Nuc. Moſch. cond. an.* ℥*ſs. Theriac. Androm.* ℥*iij. Pulv. Ari comp.* ℥*ij. cum ſ. q. Syr. Aurant. f. Elect. de quo ſumat* ℥*ij. bis in die, ſuperbibendo ſeq. Aq. diſtillat. Cochl. v. vel vj.*

℞ *Rad.*

℞ *Rad. Raphan. Ruſtic. inciſ.* ℥iij. *Cochl. hort. M. xij. Naſturt. Aq. Becabung. Salv. Menth. an. M. iv. Cort. Aurant.* N° *vj. Nuc. Moſch. cont.* N° *ij. Cerev. Brunſvicenſ.* ℔ *xij.* diſtillentur organis communibus, donec ſolummodo ℔ *vj.* aquæ eliciantur pro uſu.

Hæc autem remedia *digeſtiva*, conſtanter, atque omni adhibitâ diligentiâ, aſſiduè uſurpanda, & in *paroxyſmorum* intervallis præcipuè in uſum revocanda ſunt.

Mediocritas ea in cibo obſervanda eſt, ut neque plus alimenti ingeratur, quam ventriculus concoquere valeat, neque è contra nimiâ abſtinentiâ, partes eâ proportione alimenti defraudentur, quæ earundem robur & vigorem ſuſtinere potis eſt. Quoad qualitatem autem ciborum, ægri palato conſulendum eſt, & unica ciborum ſpecie ſingulis paſtibus veſcatur; de cæteris, demptâ carne, pro libitu comedat, modo nec acria, nec ſale condita ſint, nec aromatis. Prandere tantum expedit, loco autem cœnæ indulgeat ſibi hauſtum cereviſiæ alicujus tenuioris, quo calculi concretio impediatur.

Liquores optimi ſunt, qui neque ad vini generoſitatem aſcendunt, nec ad aquæ debilitatem, ne frigore ſuo lædant ventriculum, deprimuntve. Hujuſmodi eſt *Cereviſia tenuis Londinenſis*, aut multa aqua, cum pauco vino permixta; aqua autem pura & cruda periculoſa

loſa eſt: cum autem univerſa ferè corporis ſubſtantia, in podagram degeneraverit, à liquore quocunque fermentato quantumvis leni, & tenui abſtineat, & utatur *Decocto Diætetico ex Sarſa, in remediorum formulis deſcripto*, ubi primum à paroxyſmo convaluerit, per reliquum vitæ curſum.

Sin autem æger à longo, & nimio liquorum inebriantium uſu, vel ab ætate provectiore, vel denique præ nimia debilitate, cibos abſque vino nequeat concoquere, in paſtibus hauſtum vini Hiſpanici ſibi indulgeat, *Apozema Diæteticum* abdicans.

Æger maximè hyemis tempore, maturè lectum petat, diluculo exurgat, animus ſit tranquillus, nec nimis intentus ſtudiis, & rebus ſeriis; cæteris vero omnibus, exercitatio corporis quotidiana facile palmam præripit, quæ moderata ſit, & perſeverans, nullo autem modo juſto vehementior; quoad exercitii genus, quoties nec grandior ætas nec calculus intercedant, reliquis præferenda eſt equitatio conſtans, ac aſſidua in aere ſalubri, aut vectio in curru.

Veneris lenociniis ne indulgeat: Remedia externa nulla adhibenda.

Quamvis nihil magni in paroxyſmo tentandum ſit, conducet tamen fortaſſe per aliquot dies, ubi primum à morbo corripitur, à carnibus abſtinere, juſculo uti avenaceo, aut alio pari

pari alimento; ſin autem ſpiritus animales ataxiam concipere apti ſunt, non expedit à carnibus abſtinere diutius, quam ventriculus earum uſum adverſetur: ne quid autem vel in quantitate, vel in qualitate, ciborum aut potûs peccet.

Symptomatis iis occurrendum eſt, à quibus æger in paroxyſmo de vita periclitatur: horum maximè familiare eſt, ventriculi languor, cum ventris torminibus, tanquam à flatu: hoc in caſu, nihil æque votis reſpondet, ac vini Canarenſis hauſtulus ſubindè deglutitus, exercitio utatur æger: ſin autem ſymptoma, inducias haud ferens, mortem minitetur, modo caput malum non impetat, ad *Laudan.* ſtatim confugiendum, *viz. gutt. xx.* in hauſtulo *Aq. Epidem.* & in lecto ad quietem ſe componat: hâc ipsâ methodo, ſe ab imminenti morte liberavit, autor clariſſimus, ubi ex errore aliquo, circa res ſex non naturales, in ingentem ventriculi ægritudinem incidit, cum vomituritione, & aliquali ventris dolore, artus interim, ex retroceſſione materiæ Podagricæ, doloris erant expertes, ac ad motum plus ſolito habiles. Congium nempe unum liquoris poſſetici, vel cereviſiæ tenuis devorabat, & quamprimum totum per vomitum rejecerit, Vini Canar. hauſtulum, cui inſtillatæ fuerint *Laudan. liq. gut. xviij.* ſumebat. Quod ſi huic remedio ſymptoma non cedat, ſudor provocetur,

tur, methodo & medicamentis huic uſui dicatis, ad biduum, triduumve, manè & veſperi per duas treſve horas continuas.

Si autem metaſtaſis fuerit materiæ peccantis in Pulmonum lobos, liberatis artubus à dolore, & tumore, tractandum eſt hoc ſymptoma pari modo ac Peripneumonia abſoluta, venæſectione ſcilicet repetitâ, diætâ, & remediis refrigerantibus, & incraſſantibus; purgandus eſt inſuper æger, inter venæſectiones, potionibus lenientibus: ſudoris autem provocatio obeſt.

*FINIS.*

*Books Printed for* W. Innys *at the* Prince's Arms *in St.* Paul's *Church-yard.*

*Anthropologia Nova*: or, A new Syſtem of Anatomy, in 2 Vol. 8*vo.* with above 80 Figures drawn after the Life. By *J. Drake*, M.D.

*Pharmacopeia Extemporanea*; or, A Body of Preſcripts, by *Tho. Fuller*, M. D. 1710.

*Praxis Medica*; The Practice of Phyſick, or Dr. *Sydenham's Proceſſus Integri*, tranſlated out of *Latin* into *Engliſh* with large Annotations by *W. Salmon*, M. D. 8*vo.* 1707.

An Eſſay concerning the Knowledge and Cure of moſt Diſeaſes afflicting Human Bodies, *&c.* By *P. Paxton*, M. D. 8*vo.* 1711.

*De recta Sanguinis miſſione*; or, New and exact Obſervations on Fevers. By *J. White.* 8*vo.* 1712.

*Pharmacopeia Collegii Regalis Londini, per Ja. Shipton.* 12mo. 1711.

The Hiſtory of cold Bathing both ancient and modern, in two Parts, by Sir *J. Floyer* and Dr. *Jaynard.* 8*vo.*

*Pharmacopeia Bateana, cum Appendice Tho. Fuller, M. D.* 12mo.

*Pharmacologia, ſeu Manuductio ad Materiam Medicam, per Sam. Dale.* 2 Vol. 12mo. 1710.

Medicinal Experiments; or, A Collection of choice and ſafe Remedies, in two Parts, by *R. Boyle*, Eſq; 12*mo.* 1712.

The Anatomy of Human Bodies, with Figures drawn after the Life, and curiouſly engraven in 114 Copper Plates, on large Imperial Paper, by *W. Cowper.* Folio. Price 5*l.*